

# Week 5

Prof. K. Colaneri

Mathematics I

University of Rome Tor Vergata

11-15 October, 2021

# Outline

- ① Limits of functions
- ② Computations with limits
- ③ Undetermined forms
- ④ Intuitive and Notable limits
- ⑤ Continuity
- ⑥ Theorems on continuous functions

# Limits of functions: the intuition

Consider the following function:

$$f(x) = \frac{2x^2 - 8}{x - 2}$$

The domain is  $D = \mathbb{R} \setminus \{2\}$ .

The function is not defined in  $x = 2$ , but we can compute its value in a neighbourhood of  $x = 2$ :

| x approaching 2 from right |                 |
|----------------------------|-----------------|
| $x = 2.1$                  | $f(x) = 8.2$    |
| $x = 2.01$                 | $f(x) = 8.02$   |
| $x = 2.001$                | $f(x) = 8.002$  |
| $x = 2.0004$               | $f(x) = 8.0008$ |

| x approaching 2 from left |                 |
|---------------------------|-----------------|
| $x = 1.9$                 | $f(x) = 7.8$    |
| $x = 1.95$                | $f(x) = 7.9$    |
| $x = 1.995$               | $f(x) = 7.99$   |
| $x = 1.9991$              | $f(x) = 7.9982$ |

# Limits of functions: the intuition, cont'd

The function  $f(x)$  gets closer and closer to the value 8 as  $x$  gets closer and closer to 2.

Put in other words, if  $x$  is sufficiently close to  $s$ , then the distance between  $f(x)$  and 8 is small.

Thus, for any  $\epsilon > 0$  arbitrarily small, we can find a  $\delta > 0$  such that:

$$|f(x) - 8| < \epsilon$$

provided that

$$|x - 2| < \delta$$

# Importance of limits

Limits serve to answer the following questions:

- How does a function behave when  $x$  gets closer and closer to a point  $x_0$ ?
- How does a function behave when  $x$  gets larger and larger?

Interesting cases are particular points of the domain:

- Points where the function is not defined, i.e. **outside its domain**, but on the boundary
- $+\infty / -\infty$ , if the domain is **unbounded from above/below**

We will consider four cases:

- “Finite limit at a point”:  $\lim_{x \rightarrow x_0} f(x) = \ell$
- “Finite limit at infinity”:  $\lim_{x \rightarrow \pm\infty} f(x) = \ell$
- “Infinite limit at a point”:  $\lim_{x \rightarrow x_0} f(x) = \pm\infty$
- “Infinite limit at infinite”:  $\lim_{x \rightarrow \pm\infty} f(x) = \pm\infty$

# Finite limit at a point

**Intuitively**, we say that

$$\lim_{x \rightarrow x_0} f(x) = \ell$$


if  $f(x)$  gets close to  $\ell$  when  $x$  approaches  $x_0$

**Remark:** Approach... How?

Through points of the domain!


$$f(x) = \sqrt{x} \Rightarrow D = [0, +\infty)$$

$$f(x) = \sqrt{x} \Rightarrow D = [0, +\infty)$$






$$f(x) = \sqrt{x} \Rightarrow D = [0, +\infty)$$



$$f(x) = \frac{1}{x} \Rightarrow D = (-\infty, 0) \cup (0, +\infty)$$

$$f(x) = \frac{1}{x} \Rightarrow D = (-\infty, 0) \cup (0, +\infty)$$



$$f(x) = \frac{1}{x} \Rightarrow D = (-\infty, 0) \cup (0, +\infty)$$



We can approach  $x_0 = 0$   
from left and from right!!

# “Finite limit at a point”: $\lim_{x \rightarrow x_0} f(x) = \ell$

## Definition

Let  $f : D \subseteq \mathbb{R} \rightarrow \mathbb{R}$  be a function. Let  $x_0$  be a limit point of  $D$ . We say that


$$\lim_{x \rightarrow x_0} f(x) = \ell$$

if


$$\forall \epsilon > 0 \quad \exists \delta > 0 : \quad \forall x \in D : |x - x_0| < \delta, x \neq x_0 \Rightarrow |f(x) - \ell| < \epsilon$$

**Remark:** Note that the function is not required to be defined in  $x_0$ !


$$f(x) = \sin(x)/x$$



$$f(x) = \sin(x)/x$$



$$f(x) = \sin(x)/x$$





# Piecewise-defined functions


## Definition

A piecewise-defined function is a function defined by multiple sub-functions on different intervals.


## Example:

$$f(x) = \begin{cases} x^2 & \text{if } x < 0 \\ x & \text{if } x \geq 0 \end{cases}$$

$$f(x) = \begin{cases} x^2 & \text{if } x < 0 \\ x & \text{if } x \geq 0 \end{cases}$$



$$f(x) = \begin{cases} x^2 & \text{if } x < 0 \\ x & \text{if } x \geq 0 \end{cases}$$




# Piecewise-defined functions: the absolute value

A common example is the absolute value:

$$|x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases}$$


$$f(x) = |x|$$



# Piecewise-defined functions: other examples

Nevertheless, there is no limit in creating a piecewise-defined function...

$$f(x) = \begin{cases} x^2 & \text{if } x < 0 \\ 0 & \text{if } 0 \leq x < 1 \\ \log(x) & \text{if } x \geq 1 \end{cases}$$



# Right and left limits: the intuition

Right and left limits refer to the fact that  $x$  approaches  $x_0$  from the right or from the left.

We say that:

$$\lim_{x \rightarrow x_0^+} f(x) = \ell$$

if  $f(x)$  approaches  $\ell$  when  $x$  approaches  $x_0$  from the right, i.e.  $x > x_0$ .

We say that:

$$\lim_{x \rightarrow x_0^-} f(x) = \ell$$

if  $f(x)$  approaches  $L$  when  $x$  approaches  $x_0$  from the left, i.e.  $x < x_0$ .

# Right and left limits: the definition

## Definition

Let  $f : D \subseteq \mathbb{R} \Rightarrow \mathbb{R}$  be a function. Let  $x_0$  be a limit point of  $D$ . We say that

$$\lim_{x \rightarrow x_0^+} f(x) = \ell$$

if

$$\forall \epsilon > 0 \quad \exists \delta > 0 : \quad \forall x \in D : 0 < x - x_0 < \delta, x \neq x_0 \Rightarrow |f(x) - \ell| < \epsilon$$

similarly, we say that

$$\lim_{x \rightarrow x_0^-} f(x) = \ell$$

if

$$\forall \epsilon > 0 \quad \exists \delta > 0 : \quad \forall x \in D : -\delta < x - x_0 < 0, x \neq x_0 \Rightarrow |f(x) - \ell| < \epsilon$$



# Right and left limits: an important theorem

Why are right and left limits important?

## Theorem

*Let  $f : D \subseteq \mathbb{R} \Rightarrow \mathbb{R}$  be a function. Let  $x_0$  be a limit point of  $D$ . Then the limit*

$$\lim_{x \rightarrow x_0} f(x) = \ell$$

*exists if and only if:*

$$\lim_{x \rightarrow x_0^+} f(x) = \ell \quad \text{and} \quad \lim_{x \rightarrow x_0^-} f(x) = \ell.$$

This means that, if the right and left limits exist and are different, then we can conclude that the limit does not exist.

# Right and left limits: an example

The  $\text{sign}(x)$  function is defined on  $\mathbb{R} \setminus \{0\}$  by:

$$\text{sign}(x) = \frac{|x|}{x} = \begin{cases} -1 & \text{if } x < 0 \\ 1 & \text{if } x > 0 \end{cases}$$

If  $x > 0$ , then  $\text{sign}(x) = +1$  so that:

$$\lim_{x \rightarrow 0^+} \text{sign}(x) = +1$$


If  $x < 0$ , then  $\text{sign}(x) = -1$  so that:

$$\lim_{x \rightarrow 0^-} \text{sign}(x) = -1$$

Since the right and left limits are different, the limit does not exist:

$$\nexists \lim_{x \rightarrow 0} \text{sign}(x).$$

$$f(x) = \operatorname{sign}(x)$$



# “Finite limit at infinity”: $\lim_{x \rightarrow +\infty} f(x) = \ell$

## Definition

Let  $f : D \rightarrow \mathbb{R}$  be a function and let  $D$  be **unbounded on the right**, e.g.  $D = (a, +\infty)$  or  $D = [a, +\infty)$ . We say that

$$\lim_{x \rightarrow +\infty} f(x) = \ell$$

if

$$\forall \epsilon > 0 \quad \exists K > 0 : \quad \forall x \in D : x > K \Rightarrow |f(x) - \ell| < \epsilon$$

In this case we say that the line with equation  $y = \ell$  is an horizontal asymptote at  $+\infty$

**Important:** Notice that this case is very similar to limits of sequences. Here  $K$  plays the role as  $n^*$

# “Finite limit at infinity”: $\lim_{x \rightarrow -\infty} f(x) = \ell$

## Definition

Let  $f : D \rightarrow \mathbb{R}$  be a function and let  $D$  be **unbounded on the left**, e.g.  $D = (-\infty, b)$  or  $D = (-\infty, b]$ . We say that


$$\lim_{x \rightarrow -\infty} f(x) = \ell$$

if


$$\forall \epsilon > 0 \quad \exists K > 0 : \quad \forall x \in D : x < -K \Rightarrow |f(x) - \ell| < \epsilon$$

In this case we say that the line with equation  $y = \ell$  is an horizontal asymptote at  $-\infty$


$$f(x) = \frac{x+1}{x}$$



$$f(x) = \frac{x+1}{x}$$




$$f(x) = \frac{x+1}{x}$$






$$f(x) = \frac{x+1}{x}$$



$$f(x) = \frac{x+1}{x}$$



# “Infinite limit at a point”

It may happen that a function becomes larger and larger when  $x$  approaches a point  $x_0$ .

$$f(x) = \frac{1}{(x+5)^2}$$


The domain is  $D = \mathbb{R} \setminus \{-5\}$ . What happens when  $x$  approaches  $-5$ ?

| <hr/> x approaching $-5$ from right <hr/> |           |
|---|-----------|
| $-4.9$                                    | 100       |
| $-4.95$                                   | 400       |
| $-4.99$                                   | 1000      |
| $-4.999$                                  | 100000    |
| $-4.9999$                                 | 100000000 |

| <hr/> x approaching $-5$ from left <hr/> |           |
|--|-----------|
| $-5.1$                                   | 100       |
| $-5.05$                                  | 400       |
| $-5.005$                                 | 40000     |
| $-5.0005$                                | 4000000   |
| $-5.0001$                                | 100000000 |

$$f(x) = \frac{1}{(x+5)^2} \Rightarrow D = (-\infty, -5) \cup (-5, +\infty)$$

$$f(x) = \frac{1}{(x+5)^2} \Rightarrow D = (-\infty, -5) \cup (-5, +\infty)$$



# “Infinite limit at a point”

We now consider the case  $\lim_{x \rightarrow x_0} f(x) = \pm\infty$

## Definition

Let  $f : D \subseteq \mathbb{R} \rightarrow \mathbb{R}$  be a function and let  $x_0$  be a limit point of  $D$ . We say that


$$\lim_{x \rightarrow x_0} f(x) = +\infty$$

if


$$\forall M > 0 \quad \exists \delta > 0 : \quad \forall x \in D : |x - x_0| < \delta, x \neq x_0 \Rightarrow f(x) > M$$

In this case we say that the line with equation  $x = x_0$  is a vertical asymptote

$$f(x) = \frac{x^2 + 1}{x^2}$$




$$f(x) = \frac{x^2 + 1}{x^2}$$






$$f(x) = \frac{x^2 + 1}{x^2}$$



$$f(x) = \frac{x^2 + 1}{x^2}$$



# “Infinite limit at a point”

## Definition

Let  $f : D \subseteq \mathbb{R} \rightarrow \mathbb{R}$  be a function and let  $x_0$  be a limit point of  $D$ . We say that


$$\lim_{x \rightarrow x_0} f(x) = -\infty$$

if

$$\forall M > 0 \quad \exists \delta > 0 : \quad \forall x \in D : |x - x_0| < \delta, x \neq x_0 \Rightarrow f(x) < -M$$

In this case we say that the line with equation  $x = x_0$  is a vertical asymptote

$$f(x) = -\frac{x^2 + 1}{x^2}$$



# “Infinite limit at a point”: right and left limits

The concept of right and left limits can be extended to the case of “infinite limit at a point”.

## Theorem

Let  $f : D \subseteq \mathbb{R} \Rightarrow \mathbb{R}$  be a function. Let  $x_0$  be a limit point of  $D$ . We say that

$$\lim_{x \rightarrow x_0} f(x) = +\infty$$

**if and only if:**  $\lim_{x \rightarrow x_0^+} f(x) = +\infty$  **and**  $\lim_{x \rightarrow x_0^-} f(x) = +\infty$ . We also say that


$$\lim_{x \rightarrow x_0} f(x) = -\infty$$

**if and only if:**  $\lim_{x \rightarrow x_0^+} f(x) = -\infty$  **and**  $\lim_{x \rightarrow x_0^-} f(x) = -\infty$ .

**Remark:** If the limit does not exist (i.e. left and right limits are different), but at least one between the right and left limits are  $\pm\infty$ , **we still say that the function has a vertical asymptote at  $x = x_0$ .**

# “Infinite limit at a point”: right and left limits

$$f(x) = \frac{1}{x} \Rightarrow D = (-\infty, 0) \cup (0, +\infty)$$



We have:  $\lim_{x \rightarrow 0^-} \frac{1}{x} = -\infty$  and  $\lim_{x \rightarrow 0^+} \frac{1}{x} = +\infty$ . Therefore, the limit does not exist, however (we will see this later!) we say that the function has a vertical asymptote at  $x = 0$ .

# Infinite limit at infinity

This case covers four sub-cases:

①  $\lim_{x \rightarrow +\infty} f(x) = +\infty$

②  $\lim_{x \rightarrow -\infty} f(x) = -\infty$

③  $\lim_{x \rightarrow +\infty} f(x) = -\infty$

④  $\lim_{x \rightarrow -\infty} f(x) = +\infty$

Definitions are very intuitive. For instance  $\lim_{x \rightarrow +\infty} f(x) = +\infty$


## Definition

Let  $f : D \rightarrow \mathbb{R}$  be a function and let  $D$  be unbounded from above. We say that:  $\lim_{x \rightarrow +\infty} f(x) = +\infty$  if

$$\forall M > 0 \quad \exists K > 0 : \quad \forall x \in D : x > K \Rightarrow f(x) > M$$


**Exercise:** Write the definitions of limits for the other three cases.

$$f(x) = \frac{x^3}{10}$$







$$f(x) = \frac{x^3}{10}$$




$$f(x) = \frac{x^3}{10}$$



$$f(x) = \frac{x^3}{10}$$



$$f(x) = \frac{x^3}{10}$$



# Computations with limits

Let  $f$  and  $g$  be functions such that

$$\lim_{x \rightarrow x_0} f(x) = l, \quad \lim_{x \rightarrow x_0} g(x) = m$$

with  $l$  and  $m$  both **finite**. Then

$$\lim_{x \rightarrow x_0} f(x) + g(x) = l + m, \quad \lim_{x \rightarrow x_0} f(x) - g(x) = l - m$$

$$\lim_{x \rightarrow x_0} f(x) \cdot g(x) = l \cdot m$$

$$\text{if } m \neq 0, \quad \lim_{x \rightarrow x_0} \frac{f(x)}{g(x)} = \frac{l}{m}, \quad \text{if } m = 0 \text{ and } l \neq 0, \quad \lim_{x \rightarrow x_0} \frac{f(x)}{g(x)} = \infty$$

$$\lim_{x \rightarrow x_0} [f(x)]^p = l^p, \quad \lim_{x \rightarrow x_0} a^{f(x)} = a^l$$

$$\text{if } l > 0, \quad \lim_{x \rightarrow x_0} [f(x)]^{g(x)} = l^m$$

# Undetermined forms

Suppose that

$$\lim_{x \rightarrow x_0} f(x) = +\infty, \quad \lim_{x \rightarrow x_0} g(x) = -\infty.$$

What is limit of the sum  $f(x) + g(x)$ ?

This is an undetermined form.

**Undetermined forms**

$$+\infty - \infty, \quad , 0 \cdot \infty, \quad \frac{\infty}{\infty}, \quad \frac{0}{0}, \quad 0^0, \quad \infty^0, \quad 1^\infty.$$

# Intuitive and Notable limits

$$\lim_{x \rightarrow +\infty} a^x = \begin{cases} +\infty, & \text{if } a > 1 \\ 0, & \text{if } 0 < a < 1. \end{cases} \quad \lim_{x \rightarrow -\infty} a^x = \begin{cases} 0, & \text{if } a > 1 \\ +\infty, & \text{if } 0 < a < 1. \end{cases}$$

$$\lim_{x \rightarrow +\infty} \log_a(x) = \begin{cases} +\infty, & \text{if } a > 1 \\ -\infty, & \text{if } 0 < a < 1. \end{cases} \quad \lim_{x \rightarrow 0^+} \log_a(x) = \begin{cases} -\infty, & \text{if } a > 1 \\ +\infty, & \text{if } 0 < a < 1. \end{cases}$$

$$\lim_{x \rightarrow +\infty} \frac{\log x}{x^p} = 0, \quad p > 0.$$

$$\lim_{x \rightarrow +\infty} \frac{x^p}{a^x} = 0, \quad p > 0, a > 1.$$

$$\lim_{x \rightarrow 0^+} x^p \log x = 0.$$

# Intuitive and Notable limits

- $f(x) = x^n$ ,  $n \in \mathbb{N}$ ,  $D = \mathbb{R}$

$$\lim_{x \rightarrow +\infty} x^n = +\infty, \quad \lim_{x \rightarrow -\infty} x^n = \begin{cases} -\infty & n \text{ odd} \\ +\infty & n \text{ even} \end{cases}$$

- $f(x) = x^{-n} = \frac{1}{x^n}$ ,  $n \in \mathbb{N}$ ,  $D = \mathbb{R} \setminus \{0\}$

$$\lim_{x \rightarrow +\infty} \frac{1}{x^n} = 0, \quad \lim_{x \rightarrow -\infty} \frac{1}{x^n} = 0$$

$$\lim_{x \rightarrow 0^-} \frac{1}{x^n} = \begin{cases} -\infty & n \text{ odd} \\ +\infty & n \text{ even} \end{cases}, \quad \lim_{x \rightarrow 0^+} \frac{1}{x^n} = +\infty$$



# Limits of powers, exponentials and logarithms

- $f(x) = x^{\frac{1}{n}} = \sqrt[n]{x}$ ,  $n \in \mathbb{N}$ ,  $D = \begin{cases} \mathbb{R} & \text{if } n \text{ is odd} \\ [0, +\infty) & \text{if } n \text{ is even} \end{cases}$

- $n$  odd

$$\lim_{x \rightarrow +\infty} \sqrt[n]{x} = +\infty, \quad \lim_{x \rightarrow -\infty} \sqrt[n]{x} = -\infty$$

- $n$  even

$$\lim_{x \rightarrow +\infty} \sqrt[n]{x} = +\infty, \quad \lim_{x \rightarrow 0^+} \sqrt[n]{x} = 0$$

- $f(x) = x^{-\frac{1}{n}} = \frac{1}{x^{\frac{1}{n}}}$ ,  $n \in \mathbb{N}$ ,  $D = \begin{cases} \mathbb{R} \setminus \{0\} & \text{if } n \text{ is odd} \\ (0, +\infty) & \text{if } n \text{ is even} \end{cases}$

- $n$  odd

$$\lim_{x \rightarrow +\infty} x^{-\frac{1}{n}} = 0, \quad \lim_{x \rightarrow -\infty} x^{-\frac{1}{n}} = 0$$

$$\lim_{x \rightarrow 0^-} x^{-\frac{1}{n}} = -\infty, \quad \lim_{x \rightarrow 0^+} x^{-\frac{1}{n}} = +\infty$$

- $n$  even

$$\lim_{x \rightarrow 0^+} x^{-\frac{1}{n}} = +\infty, \quad \lim_{x \rightarrow +\infty} x^{-\frac{1}{n}} = 0$$

# Intuitive and Notable limits

$$\lim_{x \rightarrow +\infty} \left(1 + \frac{1}{x}\right)^x = e, \quad \lim_{x \rightarrow -\infty} \left(1 + \frac{1}{x}\right)^x = e$$

$$\lim_{x \rightarrow 0} \frac{\log(1+x)}{x} = 1, \quad \lim_{x \rightarrow 0} \frac{e^x - 1}{x} = 1, \quad a > 0$$

$$\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1, \quad \lim_{x \rightarrow 0} \frac{1 - \cos x}{x^2} = \frac{1}{2}$$

# Intuitive and Notable limits

If  $\lim_{x \rightarrow \dots} g(x) = +\infty$  (or  $-\infty$ )

$$\lim_{x \rightarrow \dots} \left(1 + \frac{1}{g(x)}\right)^{g(x)} = e$$

If  $\lim_{x \rightarrow \dots} h(x) = 0$

$$\lim_{x \rightarrow \dots} \frac{\log(1 + h(x))}{h(x)} = 1, \quad \lim_{x \rightarrow \dots} \frac{e^{h(x)} - 1}{h(x)} = 1$$

$$\lim_{x \rightarrow \dots} \frac{\sin(h(x))}{h(x)} = 1, \quad \lim_{x \rightarrow \dots} \frac{1 - \cos(h(x))}{h^2(x)} = \frac{1}{2}$$

Why dots? Because what really matters is the behaviour of  $h(x)$  and  $g(x)$ !

# Exercises

Compute the following limit:

$$\lim_{x \rightarrow +\infty} \frac{x-2}{x^2-3x+2}$$

This is an indeterminate form  $\frac{\infty}{\infty}$ . We use the usual trick:

$$\frac{x-2}{x^2-3x+2} = \frac{x(1-\frac{2}{x})}{x^2(1-\frac{3}{x}+\frac{2}{x^2})} = \frac{1}{x} \frac{1-\frac{2}{x}}{1-\frac{3}{x}+\frac{2}{x^2}} \rightarrow 0$$

Now, compute the following limit:

$$\lim_{x \rightarrow 2} \frac{x-2}{x^2-3x+2}$$

This time, we have an indeterminate form  $\frac{0}{0}$ . To solve these kinds of limit, observe that the equation  $x^2-3x+2=0$  has solutions  $x_1=2$ ,  $x_2=1$ . We can thus write:

$$\lim_{x \rightarrow 2} \frac{x-2}{x^2-3x+2} = \lim_{x \rightarrow 2} \frac{x-2}{(x-2)(x-1)} = \lim_{x \rightarrow 2} \frac{1}{x-1} = 1$$

# Exercises

Compute the following limit:

$$\lim_{x \rightarrow +2} \frac{x^2 - 4}{x^2 - 3x + 2}$$

It is of the form  $\frac{0}{0}$ . Factorize both the numerator and the denominator:

$$\lim_{x \rightarrow +2} \frac{x^2 - 4}{x^2 - 3x + 2} = \lim_{x \rightarrow +2} \frac{(x - 2)(x + 2)}{(x - 2)(x - 1)} = \lim_{x \rightarrow +2} \frac{x + 2}{x - 1} = 4$$

# Exercises

Compute the following limit:

$$\lim_{x \rightarrow +\infty} \sqrt{x-2} - \sqrt{x}$$

It is of the form  $+\infty - \infty$ . Multiply and divide by  $\sqrt{x-2} + \sqrt{x}$ :

$$\begin{aligned} \lim_{x \rightarrow +\infty} \sqrt{x-2} - \sqrt{x} &= \lim_{x \rightarrow +\infty} \frac{(\sqrt{x-2} - \sqrt{x})(\sqrt{x-2} + \sqrt{x})}{\sqrt{x-2} + \sqrt{x}} = \\ &= \lim_{x \rightarrow +\infty} \frac{x-2-x}{\sqrt{x-2} + \sqrt{x}} = \\ &= \lim_{x \rightarrow +\infty} \frac{-2}{\sqrt{x-2} + \sqrt{x}} = 0 \end{aligned}$$

# Exercises

Compute the following limit:

$$\lim_{x \rightarrow 0} \frac{\sqrt{1+2x} - 1}{3x}$$

It is of the form  $\frac{0}{0}$ . Multiply and divide by  $\sqrt{1+2x} + 1$ :

$$\begin{aligned} \lim_{x \rightarrow 0} \frac{\sqrt{1+2x} - 1}{3x} &= \lim_{x \rightarrow 0} \frac{(\sqrt{1+2x} - 1)(\sqrt{1+2x} + 1)}{3x(\sqrt{1+2x} + 1)} = \\ &= \lim_{x \rightarrow 0} \frac{1 + 2x - 1}{3x(\sqrt{1+2x} + 1)} = \\ &= \lim_{x \rightarrow 0} \frac{2x}{3x(\sqrt{1+2x} + 1)} = \\ &= \lim_{x \rightarrow 0} \frac{2}{3(\sqrt{1+2x} + 1)} = \frac{1}{3} \end{aligned}$$

# Exercises

Compute the following limit:

$$\lim_{x \rightarrow +\infty} \left(1 + \frac{1}{x}\right)^{3x} = \lim_{x \rightarrow +\infty} \left[ \left(1 + \frac{1}{x}\right)^x \right]^3 = e^3$$

Compute the following limit:

$$\lim_{x \rightarrow +\infty} \left(1 + \frac{3}{x}\right)^x$$

Let  $g(x) = \frac{x}{3}$  and notice that  $\lim_{x \rightarrow +\infty} g(x) = +\infty$  then we have:

$$\begin{aligned} \lim_{x \rightarrow +\infty} \left(1 + \frac{3}{x}\right)^x &= \lim_{x \rightarrow +\infty} \left(1 + \frac{1}{\frac{x}{3}}\right)^x = \lim_{x \rightarrow +\infty} \left(1 + \frac{1}{\frac{x}{3}}\right)^{\frac{x}{3} \cdot 3} \\ &= \lim_{x \rightarrow +\infty} \left( \left(1 + \frac{1}{\frac{x}{3}}\right)^{\frac{x}{3}} \right)^3 = e^3 \end{aligned}$$



# Exercises

Compute the following limit:

$$\lim_{x \rightarrow +\infty} \left(1 + \frac{7}{3x}\right)^{x-1}$$

Let  $g(x) = \frac{3x}{7}$  and notice that  $\lim_{x \rightarrow +\infty} g(x) = +\infty$ . Then

$$\begin{aligned} \lim_{x \rightarrow +\infty} \left(1 + \frac{7}{3x}\right)^{x-1} &= \lim_{x \rightarrow +\infty} \left(1 + \frac{1}{\frac{3x}{7}}\right)^{x-1} \\ \lim_{x \rightarrow +\infty} \left(1 + \frac{1}{\frac{3x}{7}}\right)^{\frac{3x}{7} \cdot \frac{7}{3}} \left(1 + \frac{1}{\frac{3x}{7}}\right)^{-1} &= e^{\frac{7}{3}} \end{aligned}$$

# Exercises

Compute the following limit:

$$\lim_{x \rightarrow 0} \frac{\log(1 + 3x^2)}{x^2}$$

Looks similar to the notable limit:

$$\lim \frac{\log(1 + g(x))}{g(x)} = 1, \quad \text{with } g(x) \rightarrow 0$$

Let  $g(x) = 3x^2$  and notice that if  $x \rightarrow 0$ , then also  $g(x) = 3x^2 \rightarrow 0$ .

Then we have

$$\lim_{x \rightarrow 0} \frac{\log(1 + 3x^2)}{x^2} = \lim_{x \rightarrow 0} \frac{\log(1 + 3x^2)}{3x^2} \cdot 3 = 1 \cdot 3 = 3$$

# Exercises

Compute the following limits:

$$\lim_{x \rightarrow 0} \frac{\sin(3x)}{2x}$$

Hint: set  $g(x) = 3x$

$$\lim_{x \rightarrow 0} \frac{2^{2x^3} - 1}{x^3}$$

Hint: set  $g(x) = 2x^3$

$$\lim_{x \rightarrow 0} \frac{\sin x}{x^3}$$

Hint: observe that  $\frac{\sin x}{x^3} = \frac{1}{x^2} \frac{\sin x}{x}$

$$\lim_{x \rightarrow 0} \frac{1 - \cos x}{x}$$

Hint: observe that  $\frac{1 - \cos x}{x} = x \frac{1 - \cos x}{x^2}$

# Asymptotes

## Vertical Asymptotes

- Compute the **Domain**
- Look for **vertical asymptotes** at **finite** limit points of the Domain
- Compute  $\lim_{x \rightarrow x_0^+} f(x)$  and  $\lim_{x \rightarrow x_0^-} f(x)$
- Is any of these  $+\infty$  or  $-\infty$ ?
  - YES  $\Rightarrow x = x_0$  is a vertical asymptote
  - NO  $\Rightarrow$  there is NO vertical asymptote

# Asymptotes

## Horizontal Asymptotes

- Compute the **Domain**
- Is  $+\infty$  and/or  $-\infty$  an extreme point of the domain?
- if YES:
  - Compute  $\lim_{x \rightarrow +\infty} f(x)$ . If the limit is a finite number  $\ell_1$ , then  $y = \ell_1$  is a horizontal asymptote at  $+\infty$ .
  - Compute  $\lim_{x \rightarrow -\infty} f(x)$ . If the limit is a finite number  $\ell_2$ , then  $y = \ell_2$  is a horizontal asymptote at  $-\infty$ .
- if the domain is NOT unbounded or any of the limits ( $\lim_{x \rightarrow +\infty} f(x)$  and/or  $\lim_{x \rightarrow -\infty} f(x)$ ) are  $\infty$ , then there is no horizontal asymptote (eventually only in one of the two sides).

# Limits of functions: exercises

Determine the asymptotes of the following function:

$$f(x) = \frac{5 - x^2}{x + 3}$$

First, we determine the domain:  $D = \mathbb{R} \setminus \{-3\}$ . Thus, we compute the limit in  $-3$ ,  $+\infty$  and  $-\infty$ . We have:


$$\lim_{x \rightarrow -3^+} \frac{5 - x^2}{x + 3} = \frac{5 - (-3^+)^2}{0^+} = -\frac{4}{0^+} = -\infty$$

$$\lim_{x \rightarrow -3^-} \frac{5 - x^2}{x + 3} = \frac{5 - (-3^-)^2}{0^-} = -\frac{4}{0^-} = +\infty$$

$$\lim_{x \rightarrow +\infty} \frac{5 - x^2}{x + 3} = \lim_{x \rightarrow +\infty} \frac{x^2 \left( \frac{5}{x^2} - 1 \right)}{x \left( 1 + \frac{3}{x} \right)} = \lim_{x \rightarrow +\infty} x \left( \frac{\frac{5}{x^2} - 1}{1 + \frac{3}{x}} \right) = +\infty \times (-1) = -\infty$$

$$\lim_{x \rightarrow -\infty} \frac{5 - x^2}{x + 3} = \lim_{x \rightarrow -\infty} \frac{x^2 \left( \frac{5}{x^2} - 1 \right)}{x \left( 1 + \frac{3}{x} \right)} = \lim_{x \rightarrow -\infty} x \left( \frac{\frac{5}{x^2} - 1}{1 + \frac{3}{x}} \right) = -\infty \times (-1) = +\infty$$

# Limits of functions: exercises



The function has a vertical asymptote in  $x = -3$ . The function has no horizontal asymptotes because  $\lim_{x \rightarrow \pm\infty} f(x) = \mp\infty$ .

# Limits of functions: exercises

Determine the asymptotes of the following function:

$$f(x) = \frac{\sqrt{x^2 + 5}}{x + 1}$$

The domain is  $D = \mathbb{R} \setminus \{-1\}$ . We compute the limit in  $-1$ ,  $+\infty$ ,  $-\infty$ .

$$\lim_{x \rightarrow -1^+} \frac{\sqrt{x^2 + 5}}{x + 1} = \frac{\sqrt{(-1^+)^2 + 5}}{0^+} = \frac{\sqrt{6}}{0^+} = +\infty$$


$$\lim_{x \rightarrow -1^-} \frac{\sqrt{x^2 + 5}}{x + 1} = \frac{\sqrt{(-1^-)^2 + 5}}{0^-} = \frac{\sqrt{6}}{0^-} = -\infty$$

$$\lim_{x \rightarrow +\infty} \frac{\sqrt{x^2 + 5}}{x + 1} = \lim_{x \rightarrow +\infty} \frac{\sqrt{x^2 \left(1 + \frac{5}{x^2}\right)}}{x \left(1 + \frac{1}{x}\right)} = \lim_{x \rightarrow +\infty} \frac{|x|}{x} \frac{\sqrt{1 + \frac{5}{x^2}}}{\left(1 + \frac{1}{x}\right)} = 1$$

$$\lim_{x \rightarrow -\infty} \frac{\sqrt{x^2 + 5}}{x + 1} = \lim_{x \rightarrow -\infty} \frac{\sqrt{x^2 \left(1 + \frac{5}{x^2}\right)}}{x \left(1 + \frac{1}{x}\right)} = \lim_{x \rightarrow -\infty} \frac{|x|}{x} \frac{\sqrt{1 + \frac{5}{x^2}}}{\left(1 + \frac{1}{x}\right)} = -1$$



# Limits of functions: exercises



The function has a vertical asymptote in  $x = -1$ . The function has two horizontal asymptotes in  $y = 1$  and  $y = -1$ .

# Limits of functions: exercises

Determine the asymptotes of the following function:

$$f(x) = \frac{x^2 + 1}{x^2 - 1}$$

The domain is  $D = \mathbb{R} \setminus \{-1, 1\}$ . We compute the limit in  $-1, 1, +\infty, -\infty$ .

$$\lim_{x \rightarrow -1^+} \frac{x^2 + 1}{x^2 - 1} = \frac{(-1^+)^2 + 1}{0^-} = \frac{2}{0^-} = -\infty$$

$$\lim_{x \rightarrow -1^-} \frac{x^2 + 1}{x^2 - 1} = \frac{(-1^-)^2 + 1}{0^+} = \frac{2}{0^+} = +\infty$$

$$\lim_{x \rightarrow 1^+} \frac{x^2 + 1}{x^2 - 1} = \frac{(1^+)^2 + 1}{0^+} = \frac{2}{0^+} = +\infty$$


$$\lim_{x \rightarrow 1^-} \frac{x^2 + 1}{x^2 - 1} = \frac{(1^-)^2 + 1}{0^-} = \frac{2}{0^-} = -\infty$$

# Limits of functions: exercises

$$\lim_{x \rightarrow +\infty} \frac{x^2 + 1}{x^2 - 1} = \lim_{x \rightarrow +\infty} \frac{x^2 \left(1 + \frac{1}{x^2}\right)}{x^2 \left(1 - \frac{1}{x^2}\right)} = \lim_{x \rightarrow +\infty} \frac{\left(1 + \frac{1}{x^2}\right)}{\left(1 - \frac{1}{x^2}\right)} = 1$$

$$\lim_{x \rightarrow -\infty} \frac{x^2 + 1}{x^2 - 1} = \lim_{x \rightarrow -\infty} \frac{x^2 \left(1 + \frac{1}{x^2}\right)}{x^2 \left(1 - \frac{1}{x^2}\right)} = \lim_{x \rightarrow -\infty} \frac{\left(1 + \frac{1}{x^2}\right)}{\left(1 - \frac{1}{x^2}\right)} = 1$$

# Limits of functions: exercises



The function has two vertical asymptotes in  $x = -1$  and  $x = 1$ . The function has one horizontal asymptotes in  $y = 1$ .

# Continuity: the intuition


Intuitively, a function is continuous if its graph is a “single unbroken” curve

## Examples

- $f(x) = x^2$ ,  $f(x) = \sin(x)$ ,  $f(x) = |x|$  are continuous functions
- $f(x) = \text{sign}(x)$ ,  $f(x) = \frac{1}{x}$ ,  $f(x) = \frac{\sin(x)}{x}$ ,  $f(x) = \begin{cases} x & x \leq -4 \\ -4 & -4 < x < 2 \\ 3 - x & x \geq 2 \end{cases}$   
are not continuous functions


# Continuity: the intuition

$$f(x) = x^2$$




# Continuity: the intuition

$$f(x) = \sin(x)$$



# Continuity: the intuition


$$f(x) = |x|$$






# Continuity: the intuition

$$f(x) = \text{sign}(x)$$




# Continuity: the intuition

$$f(x) = \frac{1}{x}$$




# Continuity: the intuition

$$f(x) = \begin{cases} \frac{\sin(x)}{x} & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases}$$



# Continuity: the intuition

$$f(x) = \begin{cases} x & x \leq -4 \\ -4 & -4 < x < 2 \\ 3 - x & x \geq 2 \end{cases}$$



# Continuity: the definition

## Definition (Continuity at a point)

Let  $f : D \rightarrow \mathbb{R}$  be a function and let  $x_0 \in D$ . We say that  $f$  is continuous in  $x_0$  if  $\lim_{x \rightarrow x_0} f(x)$  exists finite and:

$$\lim_{x \rightarrow x_0} f(x) = f(x_0)$$

## Definition (Continuity in an interval)

Let  $f : I \subseteq D \rightarrow \mathbb{R}$  be a function. We say that  $f$  is continuous in  $I$  if  $f$  is continuous in any point  $x \in I$

# Continuity: conditions for continuity

## Conditions for continuity

Suppose we have a function  $f : D \rightarrow \mathbb{R}$  and a point  $x_0 \in \mathbb{R}$ .

Then,  $f(x)$  is continuous in  $x_0$  if **ALL** these three conditions hold:

- ①  $x_0 \in D$
- ②  $\lim_{x \rightarrow x_0^+} f(x) = \lim_{x \rightarrow x_0^-} f(x) = L \quad L \neq \pm\infty$
- ③  $L = f(x_0)$

If at least one of the 3 conditions fails, we have a **discontinuity** in  $x_0$

# Classification of discontinuities

## Classification of discontinuities

Let  $f : D \rightarrow \mathbb{R}$  be a function.

- If:

$$\exists L_1 = \lim_{x \rightarrow x_0^+} f(x) \quad \text{and} \quad \exists L_2 = \lim_{x \rightarrow x_0^-} f(x)$$

but  $L_1 \neq L_2$ , the function  $f$  has a **jump discontinuity** in  $x_0$ .

- If:

$$\lim_{x \rightarrow x_0^+} f(x) = \lim_{x \rightarrow x_0^-} f(x) = L$$

but either  $L \neq f(x_0)$  or  $x_0 \notin D$ , the function  $f$  has a **removable discontinuity** in  $x_0$ .

- If at least one of the two limits:

$$\lim_{x \rightarrow x_0^+} f(x) \quad \text{or} \quad \lim_{x \rightarrow x_0^-} f(x)$$

is infinite or does not exist, the function  $f$  has an **essential discontinuity** in  $x_0$ .

# Theorem on composition of continuous functions

## Theorem

*Suppose that  $f$  and  $g$  are continuous in  $x_0$ .*

- Then  $f + g$ ,  $f - g$ ,  $f \cdot g$  are continuous in  $x_0$ .*
- If in addition  $g(x_0) \neq 0$ , then  $\frac{f}{g}$  is continuous in  $x_0$ .*
- If in addition  $f(x_0) > 0$ , then  $f^g$  is continuous in  $x_0$ .*
- If  $g$  is continuous in  $x_0$  and  $f$  is continuous in  $g(x_0) = y_0$ , then  $f \circ g$  is continuous in  $x_0$ .*
- If  $f$  is surjective, continuous and strictly increasing (or strictly decreasing) then  $f^{-1}$  is also continuous and strictly increasing (or strictly decreasing).*



# Maximum and minimum of a function


Intuitively, the maximum and minimum of a function  $f$  in a subset  $I$  of its domain  $D$  are the maximum and minimum values the function can reach in  $I$ .

## Definition


Let  $f : I \subseteq D \rightarrow \mathbb{R}$ . We say that  $f$  admits a maximum  $M \in \mathbb{R}$  in the interval  $I$  if:

- there is a point  $x_M \in I$  such that  $f(x_M) = M$
- for any other point  $x \in I$  it holds that  $f(x) \leq M$

# Example



# Example



# Maximum and minimum of a function

## Definition

Let  $f : I \subseteq D \rightarrow \mathbb{R}$ . We say that  $f$  admits a minimum  $m \in \mathbb{R}$  in the interval  $I$  if:

- there is a point  $x_m \in I$  such that  $f(x_m) = m$
- for any other point  $x \in I$  it holds that  $f(x) \geq m$

# Weierstrass theorem

Maxima and minima are not guaranteed to exist. The Weierstrass theorem provides **sufficient** conditions under which a function has maxima and minima in a subset of its domain.

## Theorem (Weierstrass Theorem - compact version)

*Any function  $f : D \rightarrow \mathbb{R}$  which is continuous function on a closed and bounded interval  $[a, b]$  admits a maximum and a minimum in  $[a, b]$  .*

## Theorem (Weierstrass Theorem - extended version)

*Let  $f : D \rightarrow \mathbb{R}$  be a function. If:*

- $f$  is continuous in an interval  $[a, b]$*
- $[a, b]$  is closed and bounded*

*then there is a point  $x_m \in [a, b]$  and a point  $x_M \in [a, b]$  such that  $f(x_m) = m$  is the **minimum** of  $f$  in  $[a, b]$  and  $f(x_M) = M$  is the **maximum** of  $f$  in  $[a, b]$ .*


# Weierstrass theorem

To apply Weierstrass Theorem three conditions must hold:

- $f$  is continuous in  $[a, b]$
- The interval  $[a, b]$  is closed (i.e. no round brackets)
- The interval  $[a, b]$  is bounded (i.e.  $a \neq -\infty$  and  $b \neq +\infty$ )

# Weierstrass theorem: examples


$$f(x) = x^2 \quad \text{in } [-2, 2]$$



The assumptions of the Weierstrass theorem are satisfied. Note that  $f$  has a minimum ( $m = 0$ ) for  $x = 0$  and a maximum ( $M = 4$ ) for  $x = -2$  and  $x = 2$ .

# Weierstrass theorem: examples

$$f(x) = x^2 \quad \text{in } (-2, 2)$$




The assumptions of the Weierstrass theorem are not satisfied because  $(-2, 2)$  is not closed. Note that  $f$  has a minimum ( $m = 0$ ) for  $x = 0$  but has no maxima, why?



# Weierstrass theorem: examples


$$f(x) = x^2 \quad \text{in } [0, +\infty)$$



The assumptions of the Weierstrass theorem are not satisfied because  $[0, +\infty)$  is unbounded. Note that  $f$  has a minimum ( $m = 0$ ) for  $x = 0$  but has no maxima, why?

# Weierstrass theorem: examples

$$f(x) = \begin{cases} x^2 & x \in [-2, 2] \setminus \{0\} \\ 1 & x = 0 \end{cases}$$



The assumptions of the Weierstrass theorem are not satisfied because  $f$  has a removable discontinuity in  $x = 0$ . Note that  $f$  has a maximum for  $x = -2$  and  $x = 2$  but has no minima, why?

# Existence of zeros (or Intermediate zero theorem)

For any continuous function in a closed and bounded interval which takes both positive and negative values in that interval, there exists at least one point in which the function is equal to zero.

## Theorem (Existence of zeros)

Let  $f : D \rightarrow \mathbb{R}$  be a function. If:

- $f$  is continuous in  $[a, b]$
- $[a, b]$  is a closed and bounded interval
- there are two points  $x_1, x_2 \in [a, b]$  such that:
  - $f(x_1) < 0$
  - $f(x_2) > 0$

then there exists a point  $x_1 < x_0 < x_2$  (or  $x_2 < x_0 < x_1$ ) such that  $f(x_0) = 0$ .

