

Quantitative Methods – I

A.Y. 2020-21

Practice 5

Lorenzo Cavallo

For any clarification/meeting: cavallo@istat.it

Summary of the Practice

1. Recap on probability distributions of discrete variables
2. Continuous Random Variables
3. Normal distribution
4. Standard Normal Distribution
5. Chi Square, Student's T and Fisher's F

THEME #1



**Recap on probability
distributions of discrete
variables**

Recap on Discrete Random Variables

For Binomial and Bernoulli sometimes you can find q instead of $(1-p)$

Bernoulli Probability Distribution

$$P(x_i) = p^{x_i} (1 - p)^{(1-x_i)} \quad E(X) = p \quad \text{and} \quad \text{Var}(X) = p \cdot (1 - p)$$

Binomial Random Variable

$$P(x) = \binom{n}{x} p^x (1 - p)^{(n-x)}$$

with n as total number of occurrences and x number of selected occurrences

$$E(X) = n \cdot p \quad \text{and} \quad \text{Var}(X) = n \cdot p \cdot (1 - p)$$

Poisson Probability Distribution

$$P(X = x) = \frac{\lambda^x e^{-\lambda}}{x!}$$

with λ as average number of occurrences in a given interval and x number of occurrences

$$E(X) = \text{Var}(X) = \lambda$$

Exercise 1

For each point, indicate first what probability distribution is used to solve the problem.

- a. 60% of people who purchase sports cars are men. If 10 sports car owners are randomly selected, find the probability that exactly 7 are men.
- b. The evaporation of water for a pool in California is (on average) 2 centimeters per month (30 days). Find the probability that the pool has lost 5 centimeters of water in August.
- c. 1% of the product sold by Apple have some bugs. Find the probability that for 100 products sold at most 1 have bugs.
- d. A company receives on average 3 returns of defective product per month. Find the probability that a randomly selected product sold in October could be defective.

Exercise 1

Solution

a. 60% of people who purchase sports cars are men. If 10 sports car owners are randomly selected, find the probability that exactly 7 are men.

a. We have to use the Binomial Distribution because the possible outcomes are only 2: men who **PURCHASE** sports car (event 1) and men who **NOT PURCHASE** sports car (event 0). Identify 'n' and 'x' from the problem.

n is the number of randomly selected items, that in this case is sports car owners, and is 10, and x, the number of men you are asked to "find the probability" for, that is 7

Find "p" the probability of success and "1-p" or "q" the probability of failure.

We are given p = 60%, or 0.6. therefore, the probability of failure is $1 - 0.6 = 0.4$ (40%)

$$P(X=7) = \frac{n!}{(n-x)! x!} \times p^x \times (1-p)^{n-x} = \frac{10!}{3! 7!} \times 0.6^7 \times 0.4^3 = 120 \times 0.0279936 \times 0.064 = 0.215 = 21.5\%$$

b. The evaporation of water for a pool in California is (on average) 2 centimeters per month (30 days).

Find the probability that the pool has lost 5 centimeters of water in August.

b. We have to use the Poisson Distribution because we have number of occurrence (cm of evaporation) in a given interval of time (a month).

Identify 'λ' and 'x' from the problem.

λ=2 (evaporation per month) and x=5 (evaporation in August)

$$P(X = 5) = \frac{\lambda^x e^{-\lambda}}{x!} = \frac{2^5 \times e^{-2}}{5!} = 0.3609 = 3.6\%$$

Exercise 1

Solution

- c. 1% of the product sold by Apple have some bugs. Find the probability that for 100 products sold at most 1 have bugs.
- c. We have to use the Binomial Distribution because the possible outcome are only 2: the product **has a bug** (event 1) and the product **has NOT a bug** (event 0). Identify 'n' and 'x' from the problem.

$$n=100 \quad x=1$$

The probability are: $p = 0.01$ and $(1-p)=0.99$

$$P(X=1) = \frac{n!}{(n-x)! x!} \times p^x \times (1-p)^{n-x} = \frac{100!}{99!1!} \times 0.01^1 \times 0.99^{99} = 100 \times 0.01 \times 0.3697 = 0.3697 = 36.97\%$$

- d. A company receives on average 3 returns of defective product per month. Find the probability that a randomly selected product sold in October could be defective.
- d. We have to use the Poisson Distribution because we have number of occurrence (return of products) in a given interval of time (a month). Identify 'λ' and 'x' from the problem.

$\lambda=3$ (return products per month) and $x=1$ (defective product sold in October)

$$P(X = 1) = \frac{\lambda^x e^{-\lambda}}{x!} = \frac{3^1 \times e^{-3}}{1!} = 3 \times 0,049 / 1 = 0,147 = 14,7\%$$

Exercise 2

On average a household receives 3 telemarketing phone calls per month. Find the probability that a randomly selected household receives:

- exactly 2 calls during a given month;
- exactly 1 call during a given week (assume 4 weeks each month);
- Find the expected value

Solution

To calculate the points a. b. and c. we have to use the Poisson probability distribution

$$P(X = x) = \frac{\lambda^x e^{-\lambda}}{x!}$$

where λ is the number of telemarketing calls received per month ($\lambda = 3$)

- We have to calculate $P(X = 2)$.

Using the Poisson distribution, $P(X = 2) = \frac{\lambda^x e^{-\lambda}}{x!} = \frac{3^2 e^{-3}}{2!} = 9 \cdot 0.049 / 2 = 0.2205 = 22.5\%$

- If the calls per month are 3, the calls per week are:

$$\lambda = 3 \text{ calls per month} / 4 \text{ weeks} = 0.75 \text{ calls per week}$$

So, $P(X = 1) = 0.75^1 \cdot e^{(-0.75)} / 1! = 0.75 \cdot 0.4724 = 0.3543 = 35.43\%$

- For the Poisson distribution: $E(X) = \text{Var}(X) = \lambda = 3$

THEME #2



Continuous Random Variables

Continuous Random Variables

- ✓ Normal distribution $X \sim N(\mu, \sigma^2)$
- ✓ Standard normal distribution $Z \sim N(0,1)$
- ✓ Chi-square distribution $X \sim \chi_n^2$
- ✓ t-Student distribution $X \sim t_n$
- ✓ F distribution $X \sim F_{n,m}$

Probability distributions of Continuous Variables

A continuous random variable X can assume any value over an interval.

2 characteristics (similar to discrete variables):

1. For any interval $[a, b]$, $0 \leq P(a < X < b) \leq 1$
2. The total probability of all the intervals within which X can assume a value is 1

$$\sum P(-\infty < X < +\infty) = 1$$


Normal distribution function, $N(\mu, \sigma^2)$

Most widely used continuous probability distribution

$$x \sim N(\mu, \sigma^2)$$

$$\text{Density function: } f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left\{-\frac{1}{2} \left(\frac{x-\mu}{\sigma}\right)^2\right\}$$

parameters: μ (mean), and σ (the spread)




Some examples of things that follow a Normal Distribution:

- Heights of people
- Size of items produced by a machine
- Errors in measurements
- Blood Pressure
- Test Scores

Normal distribution


- It is a bell-shaped curve
- Symmetry about the mean μ (mean = mode = median)
- The total area under the curve is 1 (or 100%)
- 50% of the area is to the left of the mean, and 50% to the right



The Standard Normal Distribution, Z , is a Normal Distribution with mean equal to 0 and standard deviation equal to 1, $Z \sim N(0,1)$


ANY Normal distribution, $X \sim N(\mu, \sigma^2)$, can be converted into a Standard Normal, $Z \sim N(0,1)$, by standardization:

$$Z = \frac{x - \mu}{\sigma}$$




Standard normal distribution, Z :

- It is a bell-shaped curve
- Symmetry about the mean μ (mean = mode = median)
- The total area under the curve is 1 (or 100%)
- 50% of the area is to the left of the mean, and 50% to the right
- Also for standard normal distribution




Standard normal distribution, Z:

- It is a bell-shaped curve
- Symmetry about the mean μ (mean = mode = median)
- The total area under the curve is 1 (or 100%)
- 50% of the area is to the left of the mean, and 50% to the right
- Also for standard normal distribution
- Approximately **68.3%** of the area is between $[\mu-\sigma=-1; \mu+\sigma=1]$




Standard normal distribution, Z:

- It is a bell-shaped curve
- Symmetry about the mean μ (mean = mode = median)
- The total area under the curve is 1 (or 100%)
- 50% of the area is to the left of the mean, and 50% to the right
- Also for standard normal distribution
- Approximately **68.3%** of the area is between $[\mu-\sigma=-1; \mu+\sigma=1]$
- Approximately **95.4%** of the area is between $[\mu-2\sigma=-2; \mu+2\sigma=2]$



Standard normal distribution, Z:

- It is a bell-shaped curve
- Symmetry about the mean μ (mean = mode = median)
- The total area under the curve is 1 (or 100%)
- 50% of the area is to the left of the mean, and 50% to the right
- Also for standard normal distribution
- Approximately **68.3%** of the area is between $[\mu-\sigma=-1; \mu+\sigma=1]$
- Approximately **95.4%** of the area is between $[\mu-2\sigma=-2; \mu+2\sigma=2]$
- Approximately **99.7%** of the area is between $[\mu-3\sigma=-3; \mu+3\sigma=3]$




Exercise 3

Z is a normally standard distributed variable with mean $\mu = 0$ and standard deviation $\sigma = 1$.
Find

- a. $P(z < 1.95)$
- b. $P(z > 2.32)$
- c. $P(1.19 < z < 2.12)$

Exercise 3


z	.00	.0105	...
-3.4	.0003	.00030003	...
-3.3	.0005	.00050004	...
-3.2	.0007	.00070006	...
.
.
.
1.9	.9713	.97199744	...
.
.
.
3.4	.9997	.99979997	...



a. $P(z < 1.95) = 0.9744 = 97.44\%$


Exercise 3

z	0.00	0.01	0.02
0.0	0.5000	0.5040	0.5080
0.1	0.5398	0.5438	0.5478
0.2	0.5793	0.5832	0.5871
0.3	0.6179	0.6217	0.6255
0.4	0.6554	0.6591	0.6628
0.5	0.6915	0.6950	0.6985
0.6	0.7257	0.7291	0.7324
0.7	0.7580	0.7611	0.7642
0.8	0.7881	0.7910	0.7939
0.9	0.8159	0.8186	0.8212
1.0	0.8413	0.8438	0.8461
1.1	0.8643	0.8665	0.8686
1.2	0.8849	0.8869	0.8888
1.3	0.9032	0.9049	0.9066
1.4	0.9192	0.9207	0.9222
1.5	0.9332	0.9345	0.9357
1.6	0.9452	0.9463	0.9474
1.7	0.9554	0.9564	0.9573
1.8	0.9641	0.9649	0.9656
1.9	0.9713	0.9719	0.9726
2.0	0.9772	0.9778	0.9783
2.1	0.9821	0.9826	0.9830
2.2	0.9861	0.9864	0.9868
2.3	0.9893	0.9896	0.9898



b. $P(z > 2.32) = 0.0102 = 1.02\%$

Exercise 3



c. $P(1.19 < z < 2.12) = 0.1000 = 10\%$

Exercise 4

X is a normally distributed variable with mean $\mu = 30$ and standard deviation $\sigma = 4$. Find

- $P(x < 40)$
- $P(x > 21)$
- $P(30 < x < 35)$

Solution

To solve the point a., b., c. we have to convert the Normal distribution, X in the Normal standard distribution, Z, by standardization:

$$z = \frac{x - \mu}{\sigma}$$

with mean $\mu = 30$ and standard deviation $\sigma = 4$.

- For $x = 40$, the z-value (or z-score) is: $z = (40 - 30) / 4 = 2.5$
Hence $P(x < 40) = P(z < 2.5) = [\text{area to the left of } 2.5]$ (from the Normal standard distribution table) = 0.9938
= 99.38%
- For $x = 21$, the z-score is: $z = (21 - 30) / 4 = -2.25$
Hence $P(x > 21) = P(z > -2.25) = [\text{total area}] - [\text{area to the left of } -2.25] = 1 - P(z < -2.25) = 1 - 0.0122 = 0.9878$
= 98.78%
- For $x = 30$: $z = (30 - 30) / 4 = 0$ and for $x = 35$: $z = (35 - 30) / 4 = 1.25$
Hence $P(30 < x < 35) = P(0 < z < 1.25) = \text{High value} - \text{Low value} = P(z < 1.25) - P(z < 0) =$
= $[\text{area to the left of } z = 1.25] - [\text{area to the left of } 0] = 0.8944 - 0.5 = 0.3944 = 39.44\%$

Exercise 5

A radar unit is used to measure speeds of cars on a motorway. The speeds are normally distributed with a mean of 90 km/hr and a standard deviation of 10 km/hr. What is the probability that a car picked at random is travelling at more than 100 km/hr?

Solution

Let X be the random variable that represents the speed of cars.

X has $\mu = 90$ and $\sigma = 10$.

We have to find the probability that x is higher than 100 or $P(x > 100)$

For $x = 100$, the z-score is: $z = (100 - 90) / 10 = 1$

$P(x > 90) = P(z > 1) = [\text{total area}] - [\text{area to the left of } z = 1] = 1 - P(z < 1) = 1 - 0.8413 = 0.1587$

The probability that a car selected at a random has a speed greater than 100 km/hr is the 15.87%

Exercise 6

For a certain type of computers, the length of time between charges of the battery is normally distributed with a mean of 50 hours and a standard deviation of 15 hours. John owns one of these computers and wants to know the probability that the length of time will be between 50 and 70 hours.

Solution

Let x be the random variable that represents the length of time.

It has a mean of 50 and a standard deviation of 15.

We have to find the probability that x is between 50 and 70 or $P(50 < x < 70)$

For $x = 50$ the z-score is: $z = (50 - 50) / 15 = 0$

For $x = 70$ the z-score is: $z = (70 - 50) / 15 = 1.33$ (rounded to 2 decimal places)

$P(50 < x < 70) = P(0 < z < 1.33) = [\text{area to the left of } z = 1.33] - [\text{area to the left of } z = 0] = P(z < 1.33) - P(z < 0) = 0.9082 - 0.5 = 0.4082.$

The probability that John's computer has a length of time between 50 and 70 hours is the 40,82%.

Exercise 7

Entry to a certain University is determined by a national test. The scores on this test are normally distributed with a mean of 500 and a standard deviation of 100. Tom wants to be admitted to this university and he knows that he must score better than at least 70% of the students who took the test. Tom takes the test and scores 585. Will he be admitted to this university?

Solution

Let X be the random variable that represents the scores.

X is normally distributed with a mean of 500 and a standard deviation of 100.

The total area under the normal curve represents the total number of students who took the test.

If we multiply the values of the areas under the curve by 100, we obtain percentages.

For $x = 585$ the z-score is: $z = (585 - 500) / 100 = 0.85$

The proportion P of students who scored below 585 is given by:

$P(z < 0.85) = [\text{area to the left of } z = 0.85] = 0.8023 = 80.23\%$

Tom scored better than 80.23% of the students who took the test and he will be admitted to this University.

Exercise 8

The time taken to assemble a car in a certain plant is a random variable having a normal distribution of 20 hours and a standard deviation of 2 hours. What is the probability that a car can be assembled at this plant in a period of time

- a) less than 19.5 hours?
- b) between 20 and 22 hours?

Solution

a) We have to calculate the probability that a car can be assembled in less than 19.8 hours or $P(x < 19.5)$

The z-score for $x=19.5$ is: $z = (19.5-20)/2$, so we have to calculate from the table of Z the probability:

$$P(z < -0.25) = 0.4013 = 40.13\%$$

b) We have to calculate the probability that a car can be assembled between 20 and 22 hours or $P(20 < x < 22)$

The z-score for $x=20$ is $z=0$ and for $x=22$ is $z=1$. So,

$$P(0 < z < 1) = P(z < 1) - P(z < 0) = 0.8413 - 0.5 = 0.3413 = 34.13\%$$


Chi square distribution, χ^2

Let $Z_1, Z_2, Z_3, \dots, Z_n$, be n i.i.d. random variables $Z_i \sim N(0,1)$

$$X = \sum_{i=1}^n Z_i^2 \sim \chi_n^2$$

where n =degrees of freedom (only parameter)

$$E(X) = n \quad \text{and} \quad V(X) = 2n$$



t-Student distribution, t_n

Let $Z \sim N(0,1)$ and $Y \sim \chi_n^2$,

$$X = \frac{Z}{\sqrt{Y/n}} \sim t_n$$

where n = degrees of freedom

- The total area under a t-student distribution curve is 1
- Distribution is bell-shaped and symmetric around $E(X) = 0$
- $V(X) = n/n-2$, for $n > 2$
- For $n > 60$, X converges to $Z \sim N(0,1)$

F distribution, $F_{n,m}$

Let $X \sim \chi_m^2$ and $Y \sim \chi_n^2$,


$$F = \frac{X/m}{Y/n} \sim F_{m,n}$$

where m = numerator's degree of freedom, n = denominator's degree of freedom (2 parameters)

The total area under F distribution curve is 1


Table of the Student's *t*-distribution

The table gives the values of $t_{\alpha;v}$ where
 $\Pr(T_v > t_{\alpha;v}) = \alpha$, with v degrees of freedom




α	0.1	0.05	0.025	0.01	0.005	0.001	0.0005
1	3.078	6.314	12.076	31.821	63.657	318.310	636.620
2	1.886	2.920	4.303	6.965	9.925	22.326	31.598
3	1.638	2.353	3.182	4.541	5.841	10.213	12.924
4	1.533	2.132	2.776	3.747	4.604	7.173	8.610
5	1.476	2.015	2.571	3.365	4.032	5.893	6.869
6	1.440	1.943	2.447	3.143	3.707	5.208	5.959
7	1.415	1.895	2.365	2.998	3.499	4.785	5.408
8	1.397	1.860	2.306	2.896	3.355	4.501	5.041
9	1.383	1.833	2.262	2.821	3.250	4.297	4.781
10	1.372	1.812	2.228	2.764	3.169	4.144	4.587
11	1.363	1.796	2.201	2.718	3.106	4.025	4.437
12	1.356	1.782	2.179	2.681	3.055	3.930	4.318
13	1.350	1.771	2.160	2.650	3.012	3.852	4.221
14	1.345	1.761	2.145	2.624	2.977	3.787	4.140
15	1.341	1.753	2.131	2.602	2.947	3.733	4.073
16	1.337	1.746	2.120	2.583	2.921	3.686	4.015
17	1.333	1.740	2.110	2.567	2.898	3.646	3.965
18	1.330	1.734	2.101	2.552	2.878	3.610	3.922
19	1.328	1.729	2.093	2.539	2.861	3.579	3.883
20	1.325	1.725	2.086	2.528	2.845	3.552	3.850
21	1.323	1.721	2.080	2.518	2.831	3.527	3.819
22	1.321	1.717	2.074	2.508	2.819	3.505	3.792
23	1.319	1.714	2.069	2.500	2.807	3.485	3.767
24	1.318	1.711	2.064	2.492	2.797	3.467	3.745
25	1.316	1.708	2.060	2.485	2.787	3.450	3.725
26	1.315	1.706	2.056	2.479	2.779	3.435	3.707
27	1.314	1.703	2.052	2.473	2.771	3.421	3.690
28	1.313	1.701	2.048	2.467	2.763	3.408	3.674
29	1.311	1.699	2.045	2.462	2.756	3.396	3.659
30	1.310	1.697	2.042	2.457	2.750	3.385	3.646
40	1.303	1.684	2.021	2.423	2.704	3.307	3.551
60	1.296	1.671	2.000	2.390	2.660	3.232	3.460
120	1.289	1.658	1.980	2.358	2.617	3.160	3.373
∞	1.282	1.645	1.960	2.326	2.576	3.090	3.291

Chi-Square Distribution Table



The shaded area is equal to α for $\chi^2 = \chi^2_{\alpha}$.

df	$\chi^2_{.995}$	$\chi^2_{.990}$	$\chi^2_{.975}$	$\chi^2_{.950}$	$\chi^2_{.900}$	$\chi^2_{.800}$	$\chi^2_{.700}$	$\chi^2_{.600}$	$\chi^2_{.500}$	$\chi^2_{.400}$
1	0.000	0.000	0.001	0.004	0.016	2.706	3.841	5.024	6.635	7.879
2	0.010	0.020	0.051	0.103	0.211	4.605	5.991	7.378	9.210	10.597
3	0.072	0.115	0.216	0.352	0.584	6.251	7.815	9.348	11.345	12.838
4	0.207	0.297	0.484	0.711	1.064	7.779	9.488	11.143	13.277	14.860
5	0.412	0.554	0.831	1.145	1.610	9.236	11.070	12.833	15.086	16.750
6	0.676	0.872	1.237	1.635	2.204	10.645	12.592	14.449	16.812	18.548
7	0.989	1.239	1.690	2.167	2.833	12.017	14.067	16.013	18.475	20.278
8	1.344	1.646	2.180	2.733	3.490	13.362	15.507	17.535	20.090	21.955
9	1.735	2.088	2.700	3.325	4.168	14.684	16.919	19.023	21.666	23.589
10	2.156	2.558	3.247	3.940	4.865	15.987	18.307	20.483	23.209	25.188
11	2.603	3.053	3.816	4.575	5.578	17.275	19.675	21.920	24.725	26.757
12	3.074	3.571	4.404	5.226	6.304	18.549	21.026	23.337	26.217	28.300
13	3.565	4.107	5.009	5.892	7.042	19.812	22.362	24.736	27.688	29.819
14	4.075	4.660	5.629	6.571	7.790	21.064	23.685	26.119	29.141	31.319
15	4.601	5.229	6.262	7.261	8.547	22.307	24.996	27.488	30.578	32.801
16	5.142	5.812	6.908	7.962	9.312	23.542	26.296	28.845	32.000	34.267
17	5.697	6.408	7.564	8.672	10.085	24.769	27.587	30.191	33.409	35.718
18	6.265	7.015	8.231	9.390	10.865	25.989	28.869	31.526	34.805	37.156
19	6.844	7.633	8.907	10.117	11.651	27.204	30.144	32.852	36.191	38.582
20	7.434	8.260	9.591	10.851	12.443	28.412	31.410	34.170	37.566	39.997
21	8.034	8.897	10.283	11.591	13.240	29.615	32.671	35.479	38.932	41.401
22	8.643	9.542	10.982	12.338	14.041	30.813	33.924	36.781	40.289	42.796
23	9.260	10.196	11.689	13.091	14.848	32.007	35.172	38.076	41.638	44.181
24	9.886	10.856	12.401	13.848	15.659	33.196	36.415	39.364	42.980	45.559
25	10.520	11.524	13.120	14.611	16.473	34.382	37.652	40.646	44.314	46.928
26	11.160	12.198	13.844	15.379	17.292	35.563	38.885	41.923	45.642	48.290
27	11.808	12.879	14.573	16.151	18.114	36.741	40.113	43.195	46.963	49.645
28	12.461	13.565	15.308	16.928	18.939	37.916	41.337	44.461	48.278	50.993
29	13.121	14.256	16.047	17.708	19.768	39.087	42.557	45.722	49.588	52.336
30	13.787	14.953	16.791	18.493	20.599	40.256	43.773	46.979	50.892	53.672
40	20.707	22.164	24.433	26.509	29.051	51.805	55.758	59.342	63.691	66.766
50	27.991	29.707	32.357	34.764	37.689	63.167	67.505	71.420	76.154	79.490
60	35.534	37.485	40.482	43.188	46.459	74.397	79.082	83.298	88.379	91.952
70	43.275	45.442	48.758	51.739	55.329	85.527	90.531	95.023	100.425	104.215
80	51.172	53.540	57.153	60.391	64.278	96.578	101.879	106.629	112.329	116.321
90	59.196	61.754	65.647	69.126	73.291	107.565	113.145	118.136	124.116	128.299
100	67.328	70.065	74.222	77.929	82.358	118.498	124.342	129.561	135.807	140.169




F - Distribution ($\alpha = 0.05$ in the Right Tail)

F - Distribution ($\alpha = 0.05$ in the Right Tail)

df ₂	df ₁	Numerator Degrees of Freedom								
		1	2	3	4	5	6	7	8	9
1		161.45	199.50	215.71	224.58	230.16	233.99	236.77	238.88	240.54
2		18.513	19.000	19.164	19.247	19.296	19.330	19.353	19.371	19.385
3		10.128	9.5521	9.2766	9.1172	9.0135	8.9406	8.8867	8.8452	8.8123
4		7.7086	9.9443	6.5914	6.3882	6.2561	6.1631	6.0942	6.0410	6.9988
5		6.6079	5.7861	5.4095	5.1922	5.0503	4.9503	4.8759	4.8183	4.7725
6		5.9874	5.1433	4.7571	4.5337	4.3874	4.2839	4.2067	4.1468	4.0990
7		5.5914	4.7374	4.3468	4.1203	3.9715	3.8660	3.7870	3.7257	3.6767
8		5.3177	4.4590	4.0662	3.8379	3.6875	3.5806	3.5005	3.4381	3.3881
9		5.1174	4.2565	3.8625	3.6331	3.4817	3.3738	3.2927	3.2296	3.1789
10		4.9646	4.1028	3.7083	3.4780	3.3258	3.2172	3.1355	3.0717	3.0204
11		4.8443	3.9823	3.5874	3.3567	3.2039	3.0946	3.0123	2.9480	2.8962
12		4.7472	3.8853	3.4903	3.2592	3.1059	2.9961	2.9134	2.8486	2.7964
13		4.6672	3.8056	3.4105	3.1791	3.0254	2.9153	2.8321	2.7669	2.7144
14		4.6001	3.7389	3.3439	3.1122	2.9582	2.8477	2.7642	2.6987	2.6458
15		4.5431	3.6823	3.2874	3.0556	2.9013	2.7905	2.7066	2.6408	2.5876
16		4.4940	3.6337	3.2389	3.0069	2.8524	2.7413	2.6572	2.5911	2.5377
17		4.4513	3.5915	3.1968	2.9647	2.8100	2.6987	2.6143	2.5480	2.4943
18		4.4139	3.5546	3.1599	2.9277	2.7729	2.6613	2.5767	2.5102	2.4563
19		4.3807	3.5219	3.1274	2.8951	2.7401	2.6283	2.5435	2.4768	2.4227
20		4.3512	3.4928	3.0984	2.8661	2.7109	2.5990	2.5140	2.4471	2.3928
21		4.3248	3.4668	3.0725	2.8401	2.6848	2.5727	2.4876	2.4205	2.3660
22		4.3009	3.4434	3.0491	2.8167	2.6613	2.5491	2.4638	2.3965	2.3419
23		4.2793	3.4221	3.0280	2.7955	2.6400	2.5277	2.4422	2.3748	2.3201
24		4.2597	3.4028	3.0088	2.7763	2.6207	2.5082	2.4226	2.3551	2.3002
25		4.2417	3.3852	2.9912	2.7587	2.6030	2.4904	2.4047	2.3371	2.2821
26		4.2252	3.3690	2.9752	2.7426	2.5868	2.4741	2.3883	2.3205	2.2655
27		4.2100	3.3541	2.9604	2.7278	2.5719	2.4591	2.3732	2.3053	2.2501
28		4.1960	3.3404	2.9467	2.7141	2.5581	2.4453	2.3593	2.2913	2.2360
29		4.1830	3.3277	2.9340	2.7014	2.5454	2.4324	2.3463	2.2783	2.2229
30		4.1709	3.3158	2.9223	2.6896	2.5336	2.4205	2.3343	2.2662	2.2107
40		4.0847	3.2317	2.8387	2.6060	2.4495	2.2490	2.1802	2.1240	
60		4.0012	3.1504	2.7581	2.5252	2.3683	2.2541	2.1665	2.0970	2.0401
120		3.9201	3.0718	2.6802	2.4472	2.2899	2.1750	2.0868	2.0164	1.9588
∞		3.8415	2.9957	2.6049	2.3719	2.2141	2.0986	2.0096	1.9384	1.8799

df ₂	df ₁	Numerator Degrees of Freedom									
		10	12	15	20	24	30	40	60	120	∞
1		241.88	243.91	245.95	248.01	249.05	250.10	251.14	252.20	253.25	254.31
2		19.396	19.413	19.429	19.446	19.454	19.462	19.471	19.479	19.487	19.496
3		8.7855	8.7446	8.7029	8.6602	8.6385	8.6166	8.5944	8.5720	8.5494	8.5264
4		5.9644	5.9117	5.8578	5.8025	5.7744	5.7459	5.7170	5.6877	5.6581	5.6281
5		4.7351	4.6777	4.6188	4.5581	4.5272	4.4957	4.4638	4.4314	4.3985	4.3650
6		4.0600	3.9999	3.9381	3.8742	3.8415	3.8082	3.7743	3.7398	3.7047	3.6689
7		3.6365	3.5747	3.5107	3.4445	3.4105	3.3758	3.3404	3.3043	3.2674	3.2298
8		3.3472	3.2839	3.2184	3.1503	3.1152	3.0794	3.0428	3.0053	2.9669	2.9276
9		3.1373	3.0729	3.0061	2.9365	2.9005	2.8637	2.8259	2.7872	2.7475	2.7067
10		2.9782	2.9130	2.8450	2.7740	2.7372	2.6996	2.6609	2.6211	2.5801	2.5379
11		2.8536	2.7876	2.7186	2.6464	2.6090	2.5705	2.5309	2.4901	2.4480	2.4045
12		2.7534	2.6866	2.6169	2.5436	2.5055	2.4663	2.4259	2.3842	2.3410	2.2962
13		2.6710	2.6037	2.5331	2.4589	2.4202	2.3803	2.3392	2.2966	2.2524	2.2064
14		2.6022	2.5342	2.4630	2.3879	2.3487	2.3082	2.2664	2.2229	2.1778	2.1307
15		2.5437	2.4753	2.4034	2.3275	2.2878	2.2468	2.2043	2.1601	2.1141	2.0658
16		2.4935	2.4247	2.3522	2.2756	2.2354	2.1938	2.1507	2.1058	2.0589	2.0096
17		2.4499	2.3807	2.3077	2.2304	2.1898	2.1477	2.1040	2.0584	2.0107	1.9604
18		2.4117	2.3421	2.2686	2.1906	2.1497	2.1071	2.0629	2.0166	1.9681	1.9168
19		2.3779	2.3080	2.2341	2.1555	2.1141	2.0712	2.0264	1.9795	1.9302	1.8780
20		2.3479	2.2776	2.2033	2.1242	2.0825	2.0391	1.9938	1.9464	1.8963	1.8432
21		2.3210	2.2504	2.1757	2.0960	2.0540	2.0102	1.9645	1.9165	1.8657	1.8117
22		2.2967	2.2258	2.1508	2.0707	2.0283	1.9842	1.9380	1.8894	1.8380	1.7831
23		2.2747	2.2036	2.1282	2.0476	2.0050	1.9605	1.9139	1.8648	1.8128	1.7570
24		2.2547	2.1834	2.1077	2.0267	1.9838	1.9390	1.8920	1.8424	1.7896	1.7330
25		2.2365	2.1649	2.0889	2.0075	1.9643	1.9192	1.8718	1.8217	1.7684	1.7110
26		2.2197	2.1479	2.0716	1.9898	1.9464	1.9010	1.8533	1.8027	1.7488	1.6906
27		2.2043	2.1323	2.0558	1.9736	1.9299	1.8842	1.8361	1.7851	1.7306	1.6717
28		2.1900	2.1179	2.0411	1.9586	1.9147	1.8687	1.8203	1.7689	1.7138	1.6541
29		2.1768	2.1045	2.0275	1.9446	1.9005	1.8543	1.8055	1.7537	1.6981	1.6376
30		2.1646	2.0921	2.0148	1.9317	1.8874	1.8409	1.7918	1.7396	1.6835	1.6223
40		2.0772	2.0035	1.9245	1.8389	1.7929	1.7444	1.6928	1.6373	1.5766	1.5089
60		1.9926	1.9174	1.8364	1.7480	1.7001	1.6491	1.5943	1.5343	1.4673	1.3893
120		1.9105	1.8337	1.7505	1.6587	1.6084	1.5543	1.4952	1.4290	1.3519	1.2539
∞		1.8307	1.7522	1.6664	1.5705	1.5173	1.4591	1.3940	1.3180	1.2214	1.0000



F - Distribution ($\alpha = 0.01$ in the Right Tail)

F - Distribution ($\alpha = 0.01$ in the Right Tail)


Denominator Degrees of Freedom	df ₂	Numerator Degrees of Freedom								
		df ₁	1	2	3	4	5	6	7	8
1		4052.2	4999.5	5403.4	5624.6	5763.6	5859.0	5928.4	5981.1	6022.5
2		98.503	99.000	99.166	99.249	99.299	99.333	99.356	99.374	99.388
3		34.116	30.817	29.457	28.710	28.237	27.911	27.672	27.489	27.345
4		21.198	18.000	16.694	15.977	15.522	15.207	14.976	14.799	14.659
5		16.258	13.274	12.060	11.392	10.967	10.672	10.456	10.289	10.158
6		13.745	10.925	9.7795	9.1483	8.7459	8.4661	8.2600	8.1017	7.9761
7		12.246	9.5466	8.4513	7.8466	7.4604	7.1914	6.9928	6.8400	6.7188
8		11.259	8.6491	7.5910	7.0061	6.6318	6.3707	6.1776	6.0289	5.9106
9		10.561	8.0215	6.9919	6.4221	6.0569	5.8018	5.6129	5.4671	5.3511
10		10.044	7.5594	6.5523	5.9943	5.6363	5.3858	5.2001	5.0567	4.9424
11		9.6460	7.2057	6.2167	5.6683	5.3160	5.0692	4.8861	4.7445	4.6315
12		9.3302	6.9266	5.9525	5.4120	5.0643	4.8206	4.6395	4.4994	4.3875
13		9.0738	6.7010	5.7394	5.2053	4.8616	4.6204	4.4410	4.3021	4.1911
14		8.8616	6.5149	5.5639	5.0354	4.6950	4.4558	4.2779	4.1399	4.0297
15		8.6831	6.3589	5.4170	4.8932	4.5556	4.3183	4.1415	4.0045	3.8948
16		8.5310	6.2262	5.2922	4.7726	4.4374	4.2016	4.0259	3.8896	3.7804
17		8.3997	6.1121	5.1850	4.6690	4.3359	4.1015	3.9267	3.7910	3.6822
18		8.2854	6.0129	5.0919	4.5790	4.2479	4.0146	3.8406	3.7054	3.5971
19		8.1849	5.9259	5.0103	4.5003	4.1708	3.9386	3.7653	3.6305	3.5225
20		8.0960	5.8489	4.9382	4.4307	4.1027	3.8714	3.6987	3.5644	3.4567
21		8.0166	5.7804	4.8740	4.3688	4.0421	3.8117	3.6396	3.5056	3.3981
22		7.9454	5.7190	4.8166	4.3134	3.9880	3.7583	3.5867	3.4530	3.3458
23		7.8811	5.6637	4.7649	4.2636	3.9392	3.7102	3.5390	3.4057	3.2986
24		7.8229	5.6136	4.7181	4.2184	3.8951	3.6667	3.4959	3.3629	3.2560
25		7.7698	5.5680	4.6755	4.1774	3.8550	3.6272	3.4568	3.3239	3.2172
26		7.7213	5.5263	4.6366	4.1400	3.8183	3.5911	3.4210	3.2884	3.1818
27		7.6767	5.4881	4.6009	4.1056	3.7848	3.5580	3.3882	3.2558	3.1494
28		7.6356	5.4529	4.5681	4.0740	3.7539	3.5276	3.3581	3.2259	3.1195
29		7.5977	5.4204	4.5378	4.0449	3.7254	3.4995	3.3303	3.1982	3.0920
30		7.5625	5.3903	4.5097	4.0179	3.6990	3.4735	3.3045	3.1726	3.0665
40		7.3141	5.1785	4.3126	3.8283	3.5138	3.2910	3.1238	2.9930	2.8876
60		7.0771	4.9774	4.1259	3.6490	3.3389	3.1187	2.9530	2.8233	2.7185
120		6.8509	4.7865	3.9491	3.4795	3.1735	2.9559	2.7918	2.6629	2.5586
∞		6.6349	4.6052	3.7816	3.3192	3.0173	2.8020	2.6393	2.5113	2.4073

Denominator Degrees of Freedom	df ₂	Numerator Degrees of Freedom									
		df ₁	10	12	15	20	24	30	40	60	120
1		6055.8	6106.3	6157.3	6208.7	6234.6	6260.6	6286.8	6313.0	6339.4	6365.9
2		99.399	99.416	99.433	99.449	99.458	99.466	99.474	99.482	99.491	99.499
3		27.229	27.052	26.872	26.690	26.598	26.505	26.411	26.316	26.221	26.125
4		14.546	14.374	14.198	14.020	13.929	13.838	13.745	13.652	13.558	13.463
5		10.051	9.8883	9.7222	9.5526	9.4665	9.3793	9.2912	9.2020	9.1118	9.0204
6		7.8741	7.7183	7.5590	7.3958	7.3127	7.2285	7.1432	7.0567	6.9690	6.8800
7		6.6201	6.4691	6.3143	6.1554	6.0743	5.9920	5.9084	5.8236	5.7373	5.6495
8		5.8143	5.6667	5.5151	5.3591	5.2793	5.1981	5.1156	5.0316	4.9461	4.8588
9		5.2565	5.1114	4.9621	4.8080	4.7290	4.6486	4.5666	4.4831	4.3978	4.3105
10		4.8491	4.7059	4.5581	4.4054	4.3269	4.2469	4.1653	4.0819	3.9965	3.9090
11		4.5393	4.3974	4.2509	4.0990	4.0209	3.9411	3.8596	3.7761	3.6904	3.6024
12		4.2961	4.1553	4.0096	3.8584	3.7805	3.7008	3.6192	3.5355	3.4494	3.3608
13		4.1003	3.9603	3.8154	3.6646	3.5868	3.5070	3.4253	3.3413	3.2548	3.1654
14		3.9394	3.8001	3.6557	3.5052	3.4274	3.3476	3.2656	3.1813	3.0942	3.0040
15		3.8049	3.6662	3.5222	3.3719	3.2940	3.2141	3.1319	3.0471	2.9595	2.8684
16		3.6909	3.5527	3.4089	3.2587	3.1808	3.1007	3.0182	2.9330	2.8447	2.7528
17		3.5931	3.4552	3.3117	3.1615	3.0835	3.0032	2.9205	2.8348	2.7459	2.6530
18		3.5082	3.3706	3.2273	3.0771	2.9990	2.9185	2.8354	2.7493	2.6597	2.5660
19		3.4338	3.2965	3.1533	3.0031	2.9249	2.8442	2.7608	2.6742	2.5839	2.4893
20		3.3682	3.2311	3.0880	2.9377	2.8594	2.7785	2.6947	2.6077	2.5168	2.4212
21		3.3098	3.1730	3.0300	2.8796	2.8010	2.7200	2.6359	2.5484	2.4568	2.3603
22		3.2576	3.1209	2.9779	2.8274	2.7488	2.6675	2.5831	2.4951	2.4029	2.3055
23		3.2106	3.0740	2.9311	2.7805	2.7017	2.6202	2.5355	2.4471	2.3542	2.2558
24		3.1681	3.0316	2.8887	2.7380	2.6591	2.5773	2.4923	2.4035	2.3100	2.2107
25		3.1294	2.9931	2.8502	2.6993	2.6203	2.5383	2.4530	2.3637	2.2696	2.1694
26		3.0941	2.9578	2.8150	2.6640	2.5848	2.5026	2.4170	2.3273	2.2325	2.1315
27		3.0618	2.9256	2.7827	2.6316	2.5522	2.4699	2.3840	2.2938	2.1985	2.0965
28		3.0320	2.8959	2.7530	2.6017	2.5223	2.4397	2.3535	2.2629	2.1670	2.0642
29		3.0045	2.8685	2.7256	2.5742	2.4946	2.4118	2.3253	2.2344	2.1379	2.0342
30		2.9791	2.8431	2.7002	2.5487	2.4689	2.3860	2.2992	2.2079	2.1108	2.0062
40		2.8005	2.6648	2.5216	2.3689	2.2880	2.2034	2.1142	2.0194	1.9172	1.8047
60		2.6318	2.4961	2.3523	2.1978	2.1154	2.0285	1.9360	1.8363	1.7263	1.6006
120		2.4721	2.3363	2.1915	2.0346	1.9500	1.8600	1.7628	1.6557	1.5330	1.3805
∞		2.3209	2.1847	2.0385	1.8783	1.7908	1.6964	1.5923	1.4730	1.3246	1.0000


Exercise 9

- Find the value of t for 24 df and 0.05 area in the right tail (*one-sided*);
- Find the value of t for 24 df and 0.05 in the *two-sided* area ;
- Find the probability that a r.v. t for 24 df assumes a value greater than 2.5;
- Find the probability that a r.v. t for 24 df assumes a value greater than 1.5;
- Find the probability that a r.v. χ^2 for 3 df assumes a value greater than or equal to 7;
- Find the probability that a r.v. χ^2 for 5 df leaves in the right tail a probability of 0.1;
- Find the probability that a r.v. $F_{10,10}$ leaves in the right tail a probability of 0.05.


Student's T



χ^2



F



Exercise 9

Solution

- a. We have to find the 95% percentile of the t di Student with 24 df : $t_{24,0.05}$. From the table the value is: 1.711
- b. We have to find the 2.5% e 97.5% percentile that border the two-sided area that leaves in the tail a probability of 5%. From the table $t_{24,0.025} = 2.064$ and $t_{24,0.975} = -2.064$
- c. If $X \sim t_{24}$ then, $P(X > 2.5) = 0.01$
- d. If $X \sim t_{24}$ then, $0.05 \leq P(X > 1.5) \leq 0.1$
- e. If $X \sim \chi^2_3$, then $0.05 \leq P(X > 7) \leq 0.1$
- f. From the table the 90% percentile of the distribution of the r.v. $\chi^2_{(5, 0.1)}$ with 5 df and a probability of 0.1 on the right tail is 9.236
- g. From the table the 95% percentile of the distribution of the r.v. $F_{10,10}$ and a probability on the right tail of 0.05 is 2.9782

In Teams under “General” (Generale) - “Assignment” (“Attività”) you can find the test:
“Quantitative Methods I - Test”.

The test is for you and it is used to better understand if all the themes of the first part of the course are clear.

The result of the test will not be used for the final exam.
