MATHEMATICS PRE-COURSE

Master of Science in European Economy And Business Law

Berkan Acar

berkanacar@gmail.com

SYLLABUS

Lecture 1: One variable calculus: foundations. 30 August 10AM - 1PM

Lecture 2: One variable calculus: applications. 30 August 2PM - 5PM

Lecture 3: Exponential and logarithmic Functions 31 August 10AM - 1PM

Lecture 4: Linear algebra31 August2PM - 5PM

Lecture 5: Functions of several variables. 1 September 2PM - 5PM

Reference: Mathematics for Economists, C. Simon-L. Blume. W.W. Norton & Company, Inc. ISBN 0-393-95733-0

1.One Variable Calculus Foundations $y = m(x) + 3x^{2}$ **1.1 Some Definitions**

- A real number is a value of a continuous quantity that can represent a distance along a line. It can be rational (like 3/4 or -232) or irrational (like $\pi \approx 3.14159265...$ or $\sqrt{2} \approx 1.41421356...$)
- A function (f(x)) of a real variable x with domain D is a rule that assigns a unique real number to $2\times$ $\chi \rightarrow f(\chi) = y$ each number x in D. 32-1 y=f(x) $x \xrightarrow{f} y$ x: independent(exogeneous) variable y:dependent (endogenous) variable
- The *domain* is the set of numbers x at which f(x) is defined. When the domain is not specified, it is assumed that it includes all the real numbers for which the function takes for which the function takes $(R \ 5$ meaningful values. $\rightarrow D = \mathbb{R} \setminus \{-1, 1\}$

For example for $\frac{1}{x-5}$ the domain is \mathbb{R} excluding 5. $x^2 - 1$

• The range (or co-domain) of a function is the set of all the possible values of it. FGX) e.g. for |x| the domain is $\mathbb R$ — but the co-domain is $\mathbb R^+$ **۱**×۱ wesolu

y=axn+bon-1 ____t 1.2 Function Types 5 • Polynomials : Obtained by the addition of monomials $y = ax^k$ like $h(x) = 3x^5 - 2x^2 + x + 3$ $g(x) = 3x^5 - 2x^2 + x + 3$ $2x^2 - x$ -> 3x6+4x+1 The highest exponent defines the order of the polynomial. -Constant function is polynomial of order zero: y = a-Linear functions is polynomial of order one: y = mx + n-Quadratic function (parabola) is a polynomial of order two: $y = \underline{mx^2} + nx + o$ 2x2-1x+1 -Power function is a monomial of order k: $y = ax^k$ 3x2+6x+9 • Rational functions : Ratios of polynomials $f(x) = \frac{h(x)}{g(x)} = \frac{2x+1}{3x+9}$ -A simple example is Hyperbola (constant over a monomial of order one) : y = a/xF(1) - 2 • Exponential functions : $f(\underline{x}) = e^x$ or $f(\underline{m}) = 10^{-m}$ • Trigonometric functions : $f(x) = \sin(x)$ or $f(y) = \cos(y)$ COSQ-X land= 7 Z CofQ-X sind z X

2

+

1.3 Basic geometric properties of a function

increasing or decreasing

f is increasing if $x_1 > x_2$ implies $f(x_1) > f(x_2)$ f is decreasing if $x_1 > x_2$ implies $f(x_2) > f(x_1)$

the location of its local and global minimum and maximum (if exists) ۲

The point where the function turns from decreasing to increasing is a minimum for the function

The point where the function turns from increasing to decreasing is a maximum for the function yenrete

f(x) = 2x + 1 is always increasing

f(x) = -2x + 3 is always decreasing

 $f(x) = 3x^4$ before 0 decreasing after zero increasing so it has a minimum at x=0

0

 $f(x) = -2x^2$ before 0 increasing after zero decreasing so it has a maximum at x=0

(LA)-2xt1 _x +2 92 1.4 Linear Functions Lyz 5x+2 agtb 7-5743 if y is distance, x is hours m denotes the velocity; = mx + nif y is utility, x is income m denotes marginal utility of income • for x = 0 y = n is the y-intercept, for y = 0 $x = \frac{-n}{m}$ is the x-increase of the transformation of transform ×2-×1 • Knowing two points (x_1, y_1) and (x_2, y_2) the unknowns m and n for the function can easily be found m25-1 by: ->(0, 1) $m = \frac{y_2 - y_1}{x_2 - x_1}$ substituting one point $n = y_1 - mx_1$ or $n = y_2 - mx_2$ **Exercise 1** If $0 \circ C$ equals to $32 \circ F$ and $100 \circ C$ equals to $212 \circ F$, find $\circ C$ as a linear function of $\circ F$. -0+n(X2260) 100-9212 y= 2x+1 - 180 m= 212 100 - 100 = 1.8-> F= 1.8C+37



 $f'(x_0) = \lim_{x \to 0}$

The slope of a nonlinear function of f at point $(x_0, f(x_0))$ is the slope of a nonlinear function of f at point $(x_0, f(x_0))$ is the slope of the tangent line to its graph at that $\underbrace{f(x_0+h) - f(x_0)}_{h} = \frac{df}{dx}(x_0)$

Example 1 The slope of $f(x) = x^2$ at x = 3:

$$f'(3) = \lim_{h \to 0} \frac{f(3+h) - f(3)}{h} = \lim_{h \to 0} \frac{(3+h)^2 - 3^2}{h} = \lim_{h \to 0} \frac{9 + 6h + h^2 - 9}{h} = \lim_{h \to 0} 6 + h = 6$$

YO YGt

42-YJ



7F 79 (f(x),g(x))' = f'g + f.g'(-1)' = 0 $(x-1)'_{-1}$ $(x-1) \cdot (x^{2}+x+1)' = 1 \cdot (x^{2}+x+1) + (x-1) \cdot (2x+1)$ $= x^{2} + [x + 2x^{2} + x - 2x - 1] - (x^{2} + x + 1)^{2} =$ $f(x) = 3x^{2}$ $f(3x^{2}) = \frac{1}{5}$ $f(4) = \frac{1}{5}$ $f'g - fg\overline{F}3x^2$ $\binom{7}{x-1} = \frac{5^2}{1.(x+1)}$ 1.(x+1) - (x-1).1 $\frac{x+1-x+1}{(x+1)^2} = \frac{2}{(x+1)^2}$ $(x+1)^2$ Gja f'(g(x)), g'(x) $f(f(g(x)) = (f(3x^2)) =$ $f'(3x^2)$. 6>

1.7 Differentiability

- If f(x) has a tangent line at point x_0 or $\lim_{h\to 0} \frac{f(x_0+h)-f(x_0)}{h}$ exists then f(x) is differtiable at point x_0 .
- If a function is differentiable at every point in its domain we say that it is a differentiable (smooth) function.



y2





1.10 Approximation by Differentials
Without limit
$$\frac{f(x_0+h)-f(x_0)}{h} \approx f'(x_0)$$
 or $f(x_0+h) \approx f(x_0) + f'(x_0).h$
Example 13 Assume a firm's production function is $f(x) = \frac{1}{2}\sqrt{x}$ Suppose the firm is currently using
100 unit of labor input x . Find the firms current marginal product of labor that is the additional output
that can be achieved by hiring one more unit of blom
actual result is $f'(100) = \frac{1}{4}100^{-\frac{1}{2}} = 0.025$
approximate result is $\frac{f(101)-f(100)}{1} = 0.02494$
 $f'(x) = \frac{f'(x)}{1} + \frac{f'(x)}{100} + \frac{1}{300}$
We need to find $f(1001.5) \approx f(100) + f'(1000).1.5 = 10 + \frac{1.5}{300} = 10.005$
 $f(x) = x^{1/3}$ and hence $f'(x) = \frac{1}{3}x^{-2/3}$. We know $f(1000) = 10$ and $f'(100) = \frac{1}{300}$
We need to find $f(1001.5) \approx f(100) + f'(1000).1.5 = 10 + \frac{1.5}{300} = 10.005$
 $f'(x) = \frac{8}{(t+2)^2} \rightarrow f'(0) = 2$ — the population rise $\approx f'(0).0.5 = 1$

2. One Variable Calculus: Applications

 $f'(x) = (x - 3)^2$

2.1 Using first and second derivatives to sketch graphs

- 1. Find the critical (stationary) points at which f'(x) = 0 or f'(x) is not defined, order those points showing on the x axis as $(-\infty, x_1), (x_1, x_2)...(x_k, \infty)$
 - $\times \sqrt{2}$. Find the sign of the f'(x) as x goes to ∞ for (x_k, ∞) and $-\infty$. Alternate the sign for the subsequent interval if the critical point is not even times repeated (double, quadruple etc.). Otherwise do not change the sign.
- 3. The function f(x) is increasing in the intervals with positive sign whereas it is decreasing in the intervals with the negative sign. If f'(x) is positive (negative) always then f(x) is always increasing (or decreasing).
 - 4. Do the first two steps for f''(x) instead of f'(x).

5. The function f(x) is convex (upward curved, the slope of f'(x) is increasing) in the intervals with positive sign, whereas it is concave (downward curved, the slope of f'(x) is decreasing) in the intervals with the negative sign. If f''(x) is positive (negative) always then f(x) is always convex (or concave). Positive f'' regulate f''postàlue f'

 $f(x) = \bigotimes x^2 + 2x + 1$ $F(x) = 1 \quad y = M + ercept$ $f(x) = 0 \rightarrow x = -1$



² Inflection points are the points where the function changes its concavity. They can be stationary but cannot be local minima or maxima

Example 17 Sketch $f(x) = x^4 - 8x^3 + 18x^2 - 11$ $x^2 - 6x + 9 \in (x - 6)^2 = a^2 - 2b + 6^2$

- 1. $f'(x) = 4x^3 24x^2 + 36x = 4x(x-3)^2$. Its critical points are 0 and 3. (x-3)(x-3)(x-3)
- 2. f'(x) is positive as x goes to ∞ and negative as x goes to $-\infty$, positive in $(0,\infty)$ excluding x = 3 (double root)
- 3. Then f(x) decreasing in $(-\infty,0)$ and increasing in $(0,\infty)$ excluding x=3, hence x=0 is the global minimum.

min

Saddle point

- 4. $f''(x) = 12x^2 48x + 36 = \underline{12}(x-1)(x-3) \rightarrow \text{critical points are 1 and 3}, f''(x)$ is positive as x goes to ∞ or $-\infty$
- 5. f(x) is convex in $(-\infty,1)$ and $(3,\infty)$, it is concave in (1,3).

x = 1 and x = 3 are the inflection points due to concavity changes, x = 3 is also a saddle point³ as it is a stationary (f'(3) = 0) but not an extremum point.

³ Saddle point is stationary point such that the curve (1D)/surface(2D) etc. in the neighbourhood of that point is not entirely on any side of the tangent space at that point. In one dimension, a saddle point is a stationary inflection point.

$\frac{x^{k}}{x^{m}} = \frac{x^{k}}{k^{m}} \frac{x^{k}}{k$

- In addition, we need to identify the vertical and horizontal asymtotes of the function.
- The points (x) that make the denominator of the function zero are the **vertical asymtotes.**
- If f(x) becomes close to a finite number in the limit as x goes to ∞ or $-\infty$, than that y axis point is called **horizontal asymtote**. $\int \sigma \left(|\chi| \rightarrow \langle \zeta_{\chi} \rangle \rightarrow \langle \zeta_{\chi}$

Vertical Asymptote

X=1

• For polynomials, the leading term -the monomial with the highest degree (let $a_0 x^k$)- determines the shape of the tail of the graph whether to go ∞ , $-\infty$ or an horizontal asymptote.

-if k is even both tails go to ∞ as $|x| \to \infty$ if $a_0 > 0$, both tails go to $-\infty$ as $|x| \to \infty$ if $a_0 < 0$ $(2 \times 6 + 1)$

-if k is odd one tails go to ∞ and the other goes to $-\infty$ as $|x| \to \infty$ depending on the sign of a_0

$$g(x) = \underbrace{\begin{pmatrix} a_0 x^k + a_1 x^{k-1} + \dots + a_{k-1} x^1 + a_k \\ b_0 x^m + b_1 x^{m-1} + \dots + b_{m-1} x^1 + b_m \end{pmatrix}}_{b_0 x^m} \longrightarrow \underbrace{l(x)}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0 x^m \end{pmatrix}}_{b_0 x^m} = \underbrace{\begin{pmatrix} a_0 x^k \\ b_0$$

-if k>m l(x) is a monomial, the tails of the rational function goes to $\pm \infty$ as stated above

-if k<m $l(x) \to 0$ Both the tails of g(x) are asymptotic to the x axis (y = 0) that is a horizontal asymptote for g(x)

-if k=m $l(x) \to \frac{a_0}{b_0}$ Both the tails of g(x) are asymptotic to the horizontal line $(y = \frac{a_0}{b_0})$

Example 19 Sketch
$$f(x) = \frac{1}{x}$$

 $[x^{1} \rightarrow 6 \rightarrow f(x) = y = a$ horizondal

f(x) has horizontal asymptote to the x axis. Its denominator becomes zero for x = 0 (vertical asymptote) $f'(x) = -\frac{1}{x^2}$ has a critical point at x = 0, so f(x) is decreasing in both sides of the point. $f''(x) = \frac{2}{x^3}$ is positive as x goes to ∞ and negative as x goes to $-\infty$, hence f(x) is convex on the right and concave on the left of y axis.





2.3 Maxima and Minima

• Finding the maximum or minimum points (extremum points) is very important in economics in order to reach optimal values of economic variables. e.g. maximizing utility or profit, minimizing cost.

2

-> 0t

RA

6,

• An extremum point can be on the boundary or in the interior part of the domain.

• If it is an interior extremum point then it is a critical point. For the critical points:
-if
$$f'(x_0) = 0$$
 and $f''(x_0) < 0$, then x_0 is a maxima of $f(x)$
-if $f'(x_0) = 0$ and $f''(x_0) > 0$, then x_0 is a minima of $f(x)$
-if $f'(x_0) = 0$ and $f''(x_0) = 0$, then x_0 can be a minima (e.g. $f(x) = x^4$), maxima (e.g. $f(x) = -x^4$) or
neither (e.g. $f(x) = x^3$).
Example 22 Find minima and maxima of $f(x) = x^4 - 4x^3 + 4x^2 + 4$
 $f'(x) = 4x^3 - 12x^2 + 8x = 4x(x-1)(x-2)$, critical points are 0.1 and 2
 $f''(x) = 12x^2 - 24x + 8$
 $f''(0) = f''(2) = 8 > 0 \rightarrow x = 0$ and $x = 2$ are minimum points $f''(1) = -4 < 0 \rightarrow x = 1$ is a maxima

Global Maxima and Minima

In three cases global maximum or minumum points are found easily:

- 4 A
- 1. When the domain of f(x) is an interval in \mathbb{R}^{-1} and it has only one critical point that is a local max (min) then that point is global max (min)

The proof comes from the idea that if the point were not a global max (min) then there should be another critical point between two max (min). $f'' \ge 0$ f_1 increasing

- 2. If f is a C² function whose domain is an interval and f'' is never zero, then f has at most one critical point that is a global min f''(x) > 0 and a global max f''(x) < 0 The proof: If f'' > 0 always then f' is an increasing function such that it can have a value "0" only one time and from the previous theorem it is global min.
- 3. A continuous functions whose domain is a closed and bounded interval [a,b] must have a global max and a global min (Weierstrass theorem).

In this case, we find critical points and evaluate the function at these points and the boundary points. The point with largest value of f is the global maximum and the point with the smallest vlue of is the global minimum.

Example 23 A firm produces every book with a cost of \$5. Every book is sold for \$10 and each day 10 books are sold currently. The firm expects to sell one additional book for each dollar decrease in price. What are the demand and profit functions and the price P that maximizes the profit?

X = mP + n, m = -1, current (X, P) = (10, 10), substituting them in the demand equation we find X = 20 - P then the profit becomes: $\pi = (P - 5)(20 - P) = -P^2 + 25P - 100$ $\pi' = -2P + 25 \rightarrow P = 12.5$ is the only critical point, $\pi'' = -2 < 0$ then P = 12.5 is the global maximum point that maximes the profit of the firm.



2.4 Applications to Economics

2.4.1 Production Function

- It relates the amount of output (x) to the amount of input (q): x = f(q)
- Continuous or maybe C^2
- Increasing
- There is a level of input until which the function is convex and then it is concave



2.4.2 Cost Function

- A cost function assigns a cost to the level of output: C(x)
- MC(x) = C'(x) is the marginal cost that measures the additional cost incurred from the production
 of one unit more when the current output is x.
- The average cost is the cost per unit produced: AC(x)=C(x)/x
- The essential properties of cost function:
 C(x) is a naturally increasing function. Moreover, Let C(x) is C¹ then
 -if MC>AC, AC is increasing (Doing better than average rises average)
 -if MC<AC, AC is decreasing (Doing worse than average decreases average)
 -at an interior minimum of AC, MC=AC

The proof comes from:
$$AC'(x) = \frac{d}{dx}\left(\frac{C(x)}{x}\right) = \frac{C'(x)x - C(x)}{x^2} = \frac{C'(x) - C(x)/x}{x} = \frac{MC - AC}{x}$$

2.4.3 Revenue and Profit Functions

- How much money a firm receives from selling its x unit of output: R(x) = p(x)x
- MR(x) = R'(x) is the marginal revenue, average revenue(AR) is the unit price p(x)
- Profit π(x) = R(x) − C(x) to maximize it we solve for π'(x*) = MR(x*) − MC(x*) = 0 → MR(x*) = MC(x*). This means optimum output (x*) for maximum profit occurs when MR = MC.
- In a model of perfect competetion (with many firms and no individual firm can control the output price by its production activity) the market price for any firm receives for its output is constant: p(x) = p R(x) = px MR = AR = p



- MC curve gives the locus of optimal price output combinations. MC curve is the firm's supply curve which relates the market price to the amount produced.
- Under perfect competetion $\pi'(x) = p C'(x) \rightarrow \pi''(x) = -C''(x)$

Profit maximizing $\pi''(x) \leq 0$ implies $C''(x) \geq 0$ At the optimal output the firm experiences increasing MC.

$$f(x) = a^{x} \quad b^{x} \quad 2^{x} \quad \Pi^{x}$$

3 Exponential and Logarithmic Functions

$$f(t) = a^{t} \quad \text{where } a > 0 \text{ is called an exponential function.}$$

$$f(t) = a^{t} \quad \text{where } a > 0 \text{ is called an exponential function.}$$

$$f(t) = a^{t} \quad \text{where } a > 0 \text{ is called an exponential function.}$$

$$f(t) = a^{t} \quad \text{where } a > 0 \text{ is called an exponential function.}$$

$$f(t) = a^{t} \quad \text{where } a > 0 \text{ is called an exponential function.}$$

$$f(t) = a^{t} \quad \text{where } a > 0 \text{ is called an exponential function.}$$

$$f(t) = 1 \text{ by definition}$$

$$f(t) = \sqrt[n]{a} \text{ nth root of a}$$

$$f(t) = \sqrt[n]{a} \text{ nth root of a}$$

$$f(t) = \sqrt[n]{a} \text{ mth power of the nth root of a}$$

$$f(t) = a^{t} = \frac{\sqrt{1}}{a^{|t|}} \text{ that is the reciprocal of } a^{|t|}$$

$$\int_{a} f^{x} = \sqrt[n]{a} f^{x}$$

$$\int_{a} f^{x} = \sqrt[n]{a} f^{$$

Example 28 Graphs of
$$2^{-t}, 5^{-t}, 8^{-t}, 2^{t}, 5^{t}$$
 and 8^{t} $(f) = 9 = a^{t}, f_{0} = 1$

Negative exponents are the mirror image of the pozitive ones with respect to y axis. As the base increases the function becomes steeper.



 $\frac{1}{\left(1+\frac{1}{2h}\right)^{2h}} \stackrel{2}{\Rightarrow} \frac{1}{\left(1+\frac{1}{2h}\right)^{n}} \stackrel{4}{\Rightarrow} \frac{1}{\left(1+\frac{1}{2h}\right)^{n}} \stackrel{4}{\wedge} \frac{1}{\left(1+\frac{1}{2h}\right)^{n}} \stackrel{4}{\wedge} \frac{1}{\left(1+\frac{1}{2h}\right)^{n}} \frac{1}{\left(1+\frac{1}{2h}\right)^{n}} \stackrel{4}{\Rightarrow} \frac{1}{$ Euler's number (e), logarithm and ln

• The number $e = \lim_{n \to \infty} \left(1 + \frac{1}{n}\right)^n \approx 2.7$

lm A. (1+ E) 52.7 If one deposits A Euro in an account which pays an annual interest rate r compounded continuously, then after t years the account will grow to Ae^{rt}

$$\begin{pmatrix} e^{y} \\ e^{z} \\ e^{z} \\ e^{z} \end{pmatrix}^{n} = c^{n} \\ \lim_{n \to \infty} \left(1 + \frac{r}{n} \right)^{n} = \lim_{n \to \infty} \left(1 + \frac{1}{\frac{n}{r}} \right)^{n} \text{ let } \frac{n}{r} = m \to \lim_{n \to \infty} \left(1 + \frac{1}{m} \right)^{rm} = e^{r} \\ \begin{pmatrix} e^{z} \\ A \\ e^{z} \\ e^{z$$

- Base 10 logarithm=Log $y = Logx \rightarrow 10^y = x$ $Log1000 = Log10^3 = 3$ Base e logarithm=ln $\ln x = y \rightarrow e^y = x$ $\ln z = 0$ $\ln z = 0$
- The graphs of e^x , $\ln x$, 10^x , Logx for which the logarithms are the mirror image of the exponential functions with the same base with respect to y = x line. As the base increases logarithm functions $, \log \sim ln$ becomes less steeper and closer to x axis as $x \to \infty$





pounded continuously?

 $2A = A.e^{+i}$ $ln(2 = e^{+i})$ lne = 1 lne = x ln et = + $\ln 2 = r + \frac{\ln 2}{r} = \frac{\ln^2}{\ln^2} = \frac{\ln^2}{0.1} \simeq 6.9 \text{ years}$ r-10%

$$(2x^{2}) = hy \quad (x^{5})^{2} fx^{h}$$
4.1 Derivatives of exponential and logarithm:

$$(e^{x})' = e^{x} \quad (hx^{h})' = hx^{h}$$

$$\stackrel{d}{dx} (\lim_{n \to \infty} (1 + \frac{r}{n})^{n}) = \lim_{n \to \infty} n (1 + \frac{r}{n})^{n-1} \frac{1}{n} = \lim_{n \to \infty} (1 + \frac{r}{n})^{n-1} = \lim_{n \to \infty} (1 + \frac{r}{n})^{n} \text{ as } n \to \infty$$

$$b)(\ln x)' = \frac{1}{x}$$

$$(\ln x)' = \lim_{h \to 0} \frac{\ln(x+h) - \ln(x)}{h} = \lim_{h \to 0} \frac{1}{h} \ln(\frac{x+h}{x}) = \lim_{h \to 0} \ln(1 + \frac{h}{x})^{\frac{1}{h}} = \lim_{h \to 0} \ln(1 + \frac{1/x}{1/h})^{\frac{1}{h}} = \ln e^{1/x} = 1/x$$

$$(\ln x)' = e^{u(x)} \cdot u'(x) \text{ obtained by chain rule}$$

$$d) (\ln(u(x))' = \frac{u'(x)}{u(x)} \text{ if } u(x) > 0 \text{ obtained by chain rule}$$

$$e) (b^{x})' = (e^{x \ln b})' \text{ using } c) \quad (b^{x})' = e^{x \ln b} \cdot \ln b = b^{x} \cdot \ln b$$

$$(e^{x}) = e^{x} \quad (b^{x})' = b \cdot (xb)$$





4.2 Applications

4.2.1 Present Value



If we put B Euros into a saving account with annual interest rate r which is compounded continuously, then after t years it becomes $A = Be^{rt}$. Conversely in order to generate A Euros t years from now in an account compunded interest rate r continuously, we would have to invest $B = Ae^{-rt}$ Euros that is the present value (**PV**) of B Euros t years from now.

Annuity is a sequence of payments at regular intervals over a specified period. The present value of an anuity that pays A Euros at the end of the next N years with an interest rate r of continous compounding. $PV = Ae^{-r} + Ae^{-2r} + ... + Ae^{-nr} = A(e^{-r} + e^{-2r} + ... + e^{-nr})$ Using the property of geometric series⁵: $= A \frac{e^{-r}(1-e^{-rn})}{1-e^{-r}} = A(e^{-rn}) e^{r-1}$

Example 33 Assuming 10% interest rate compunded continuously, what is the present value of an annuity thay pays 500 Euros a year? n = 5 r = 0.

a) for the next 5 years:
$$500(1-e^{-0.5}) = 1870.6$$

b) forever: $\frac{500}{e^{0.1}-1} = 4754.2$
 $X = a + a^1 + ... + a^n = \frac{a(1-a^n)}{1-a}$ (Found by subtracting X/a from X)



 $PV = \left(\frac{A}{1+r} + \frac{A}{(1+r)^2} + \dots + \frac{A}{(1+r)^n} \text{ using again geometric series expansion}\right)$ $= A \frac{\frac{1}{1+r}(1 - \left(\frac{1}{1+r}\right)^n)}{1 - \frac{1}{1+r}} = \frac{A}{r} \left(1 - \left(\frac{1}{1+r}\right)^n\right)$ $= \frac{A}{r} \text{ as } n \to \infty$

In order to generate A Euros per year from an account paying annual interest with rate r one must deposit inti the account $\frac{A}{r}$ initially.

Example 34 Redo the previous example with annual compunding? a) for the next 5 years: $\frac{500}{0.1} \left(1 - \left(\frac{1}{1.1}\right)^5\right) = 1895.4$ b) forever: $\frac{500}{0.1} = 5000$
4.2.2 Optimal Holding Time

Suppose the market value of your real estate will be V(t) Euros t years from now. If the interest rate remains constant (r) and continuously compunded during this period then the present value of the real estate is $V(t)e^{-rt}$. Maximizing the present value gives the optimal time to sell it.

$$\left(V(t)e^{-rt} \right)' = V'(t)e^{-rt} - rV(t)e^{-rt} = 0$$

$$\frac{V'(t)}{V(t)} = r \text{ at the optimal selling time}$$

percent growth rate of the value of the real estate =percent rate of change of money in the bank

Example 37 The value of a land is increasing according to the formula $V = 2000e^{t^{\frac{1}{4}}}$. If the interest rate is 10%, how long it should be held to max its present value?

$$\ln V = \ln 2000 + t^{\frac{1}{4}} \to (\ln V)' = \frac{V'(t)}{V(t)} = r = \frac{1}{4}t^{-\frac{3}{4}} \to r = 0.1 \text{ then } t = 3.39$$

5 Linear Algebra

In general, an equation is linear if it has the form

$$a_1x_1 + a_2x_2 + \dots + a_nx_n = b$$

where the letters
$$a_1, a_2, ..., a_n, b$$
 are the fixed numbers and therefore the are called parameters wheras $x_1, x_2, ..., x_n$ stand for the variables.

 $3x_1 + 2x_2 = 5$

4 + 22=2

As a key feature of linear equations, each term contains at most one variable and that variable can have only the first power. Linear equations are easy to handle (they build on the techniques learned in high school, such as the solution of two linear equations in two unknowns such as substitution or elimination of variables and they build on simple geometry of the plane and the cube which are easy to visualize). These equations can often have exact solutions unlike nonlinear systems. With suitable assumptions or linearizations they can be good approximations of the nonlinear systems. Moreover, some of the most frequently studied models



5.1.1 Tax benefits of charitable contributions

A firm earns before-tax profits of \$100,000. It has agreed to donate 10% percent of its after-tax profits to the a charity fund. It must pay a state tax of 5 percent of its profits (after the donation) and a federal tax of 40 percent of its profits (after the donation and state taxes are paid). How much does the company pay in state taxes, federal taxes, and charitable donation?

Xty=

- Let S, F and C are state taxes, federal taxes, and charitable donation. After tax profits are 100,000-(S+F); so C becomes $C = 0.1(100,000 (S+F)) \rightarrow C + 0.1S + 0.1F = 10,000$.
- S is 5% of profits net of the donation then $S = 0.05(100.000 C) \rightarrow 0.05C + S = 5,000$
- Federal taxes are 40% the profit after deducting C and S \rightarrow $F = 0.4(100.000 C S) \rightarrow 0.4C + 0.4S + F = 40,000$

In summary we get three linear equations, subsituting the second equation to the others we get two equations with two unknowns:

$$\frac{1}{12} = \frac{10}{10} = \frac{10}{10} = \frac{10}{10} = \frac{10}{100} = \frac{10}{10$$

Then the solution becomes C=5956 S=4702 F=35737 and after tax&contribution profit is \$53605

Without donation the after tax profits become \$57000 meaning that \$5956 donation costs \$3395 to the firm.

5.1.2 Linear Model of Production

As a simplification constant return to scale production is assumed that is the amount of output linearly proportional to the amount of input. e.g. 50 cars need 50 times the input of one car.

• In an open Leontief system of economy, the production of a good i (there are n+1 goods in the economy) can be described by a set of input output coefficients where a_{ij} denotes the input of good i needed to produce one unit of good j. The output of good i must be allocated between production activities and consumption. Good 0 is labor that is supplied by consumers so the consumption (demand) for each good i is given exogenously (this is why it is called open system) that is not solved for in the model. Each good i is used for producing other goods and consumption (c_i) . Good "0" is labor that is supplied by consumers so its consumption c_0 is negative.

$$x_i = a_{i1}x_1 + a_{i2}x_2 + \ldots + a_{in}x_n + c_i$$

- Explicitly (gross output=used as input+consumption):
 - $(1 a_{11})\underline{x_1} a_{12}x_2 \dots a_{1n}\underline{x_n} = c_1$ $-a_{21}x_1 + (1 a_{22})x_2 \dots a_{2n}x_n = c_2$

$$-a_{n1}x_1 \qquad -a_{n2}x_2 - \dots + (1 - a_{nn})x_n = c_n$$
$$-a_{01}x_1 \qquad -a_{02}x_2 - \dots - a_{0n}x_n = c_0$$

 $X_{1} = 0 X_{1} + 0.2 X_{2} + 0.5 X_{3}$

5.1.3 Markov Models of Employment

- Using transition probabilities from unemployment to employment or from employment to uneployment, these models are commonly used to understand the log run employment behaviour.
- Let x_t and y_t are the number of employed and unemployed, q and p are the probability of employment for them respectively. Assuming finding job or leaving job are independent of number of weeks worked. Then the number of employed and unemployed in the next period (say week) is given as:

$$x_{t+1} = qx_t + py_t$$

 $y_{t+1} = (1-q)x_t + (1-p)y_t$

• Normalizing the total number of employed and unemployed to 1, in the steady state:

$$x = qx + py$$

$$y = (1 - q)x + (1 - p)y$$

$$x + y = 1$$

• The first two equations are the same equations with minus sign so in fact we have two equations

• Then
$$x = \frac{p}{1+p-q}$$
 and $y = \frac{1-q}{1+p-q}$
• Using $q = 0.998$ and $p = 0.136$ (Hall,1966) for US white males in 1966
 $x = \frac{0.136}{1+0.136-0.998} = 0.986$ $y = 1 - x = 1.4\%$ of white males were unemployed on average in 1966.

5.1.4 IS-LM Analysis

IS (Investments, Savings) and LM(Liquidity, Money) analysis is a linear model of a closed economy with total national income (Y) total national spending (Consumption, Investment and Government expenditures)

- Y = C + I + GY. by z swing (1-b) y • For IS analysis, on the consumer side spending is proportional to total income C = bY (0 < b < 1) where b is called marginal propensity to consume, while s=1-b is the marginal propensity to save,
- On the firms side, either they invest to keep their money in the bank with an interest rate (r) so investment is a decreasing function of r.

 $\underbrace{Y = bY}_{\textbf{SY}} + I^0 - ar + G$

 $sY + ar = I^0 + G$

 $\gamma = (-\frac{1}{2}r - 1] + 6$

- Or we write it as
- IS equation describes the real side of the economy by summarizing consumption, investment and saving decisions.

On the other hand, the LM equation is determined by the money market equilibrium condition that money supply (M_s) equals money demand (M_d) . M_d has two components: the transactions or precautionary demand (M_{dt}) and the speculative demand (M_{ds}) . The transactions demand derives from the fact that most transactions are denominated in money. Thus, as national income rises, so does the demand for funds. So that $M_{dt} = mY$

The speculative demand comes from the portfolio management problem faced by an investor in the economy. The investor must decide whether to hold bonds or money. Money is more liquid but returns no interest, while bonds pay at rate r. It is usually argued that the speculative demand for money varies inversely with the interest rate (directly with the price of bonds). The simplest such relationship is the linear one:

$$M_{ds} = M^0 - hr$$

Equating the supply to the demand:

$$M_s = mY + M^0 - hr$$



Hence we can write IS-LM system of equations as:

$$sY + ar = I^0 + G$$
$$mY - hr = M_s - M^0$$

where the solution (Y, r) depend upon the policy parameters M_s , and G and on the behavioral parameters a, h, I^0, m, M^0 and s.

Example 38 Consider the above model with no fiscal policy (G = 0). Suppose that $M_s = M^0$: that is. the intercept of the LM curve is 0. Suppose that $I^0 = 1000, s = 0.2, h = 1500, a = 2000, and m = 0.16$. Write out the explicit IS-LM system of equations. Solve them for the equilibrium GNP Y and the interest rate r.



5.2 Systems of Linear Equations

There are essentially three ways of solving systems of linear equations: substitution, elimination of variables, and matrix methods.

5.2.1 Substitution and elimination of variables methods

- Substitution is simply made by writing one variable in terms of other(s) using an equation and substituting this relation into the other equation(s).
- Elimination of variables is generally more conducive to the theoretical analysis. It is done by multiplying equations and adding them up such that eliminating unknown(s) to solve the equation with less unknowns. This is called Gauss elimination.

Example 40 An example for linear production model (look section 5.1.2 for detail) of 3 goods (x_1, x_2, x_3) given their production input-output proportions and exogenous consumption amounts (130,74,95) can be written as:

$$\begin{array}{rcl} x_1 &=& 0x_1 + 0.4x_2 + 0.3x_3 + 130 \rightarrow x_1 - 0.4x_2 - 0.3x_3 = 130 \\ x_2 &=& 0.2x_1 + 0.12x_2 + 0.14x_3 + 74 \rightarrow -0.2x_1 + 0.88x_2 - 0.14x_3 = 74 \\ x_3 &=& 0.5x_1 + 0.2x_2 + 0.05x_3 + 95 \rightarrow -0.5x_1 - 0.2x_2 + 0.95x_3 = 95 \end{array}$$

substituting $x_1 = 0.4x_2 + 0.3x_3 + 130$ into the other equations :

$$\begin{array}{rcl} -0.2 \left(0.4x_{2} + 0.3x_{3} + 130 \right) + 0.88x_{2} - 0.14x_{3} &= 74 \\ -0.5 \left(0.4x_{2} + 0.3x_{3} + 130 \right) - 0.2x_{2} + 0.95x_{3} &= 95 \end{array} \\ \begin{array}{rcl} & & & & & & & & \\ & & & & & & & \\ & & & & & & & \\ & & & & & & & \\ & & & & & & & \\ & & & & & & & \\ & & & & & & & \\ & & & & & & & \\ & & & & & & & \\ & & & & &$$

Example 41 We do the previous example with elimination of variables. Multiplying the first one by 0.2 and adding it to second to eliminate x_1 ; multiplying the first one by 0.5 and adding it to the third to eliminate x_1 :

$$0.2(x_1 - 0.4x_2 - 0.3x_3) = 0.2 * 130$$

$$+ -0.2x_1 + 0.88x_2 - 0.14x_3 = 74$$

$$0.8x_2 - 0.2x_3 = 100$$

$$\begin{array}{r} 0.5 \left(x_1 - 0.4x_2 - 0.3x_3\right) = 0.5 * 130 \\ + 0.5x_1 - 0.2x_2 + 0.95x_3 = 95 \\ \hline -0.4x_2 + 0.8x_3 = 160 \end{array}$$
 Then multiplying the above found by 0.5 and adding it to the this:

$$0.5(0.8x_2 - 0.2x_3) = 0.5 * 100 + -0.4x_2 + 0.8x_3 = 160 0.7x_3 = 210$$

means our system transforms into

 $x_3 = 300$ by back subtitution into the others we find $x_2 = 200$ and $x_1 = 300$

.

6. Matrix Algebra

• We can write a linear system of equations in matrix form.

$$A.x = b$$

$$X = A^{-1}.b$$

- In compact form we can write as Ax = b Where the matrix A is the coefficient matrix, x vector of variables, b vector of constants
- For example:

/

• Where the matrix
$$A = \begin{bmatrix} 0.8 & -0.2 \\ -0.4 & 0.8 \end{bmatrix}$$
 is the coefficient matrix, $x = \begin{bmatrix} x_2 \\ x_3 \end{bmatrix}$ vector of variables,

$$b = \begin{bmatrix} 100 \\ 160 \end{bmatrix}$$
 vector of constants
• The size of a matrix is p, x, k, where p is the number of rows and k is the number of columns; for

• The size of a matrix is n x k where n is the number of rows and k is the number of columns; for example, a 2 x 3 is a matrix with 2 rows and 3 columns; a n x n matrix is called square matrix (same number of rows and columns)



6.1 Addition or subtraction

• In order to perform algebraic operations, matrices must meet some requirements about their size. For addition or subtraction, they have to be the same size. Each element of the matrix added to or subtracted from the element in the same position. $A\pm B$

6.2 Scalar multiplication

There is no size requirement for scalar multiplication. Each element is multipled by the scalar. \mathbf{rA}

$$r\begin{bmatrix} a_{11} & . & . & a_{1n} \\ . & . & . \\ . & a_{ij} & . \\ . & . & . \\ a_{k1} & . & . & a_{kn} \end{bmatrix} = \begin{bmatrix} ra_{11} & . & . & ra_{1n} \\ . & . & . \\ . & ra_{ij} & . \\ . & . & . \\ ra_{k1} & . & . & ra_{kn} \end{bmatrix}$$

6.3 Matrix multiplication

 $\bullet\,$ We can define the matrix product \underline{AB} if and only if

number of columns of \mathbf{A} =number of rows of $\underline{\mathbf{B}}$

- If A is k x m matrix and B is m x n matrix AB becomes k x n : (kxm)(mxn) = (kxn)
- IA = A where I = (kxk) Identity matrix (with all ones on the diagonal and other terms 0)
- To obtain the (i,j)th entry of AB, multiply the ith row of A and the jth column of B as the following:

A, D = kxm mx n A.B Kxr

 $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

6.5 Laws of Matrix Algebra

6x2:2x3 x 2x3 4x2

- Associative: (A+B)+C=A+(B+C) (AB)C=A(BC)
- Commutative for addition: A+B=B+A not for multiplication $AB\neq BA$
- Distributive Laws: A(B+C)=AB+AC (A+B)C=AC+BC
- Transpose (writing rows as columns and columns as rows). If A is k x n its transpose A^{T} (or A') is n x k 2 x 2

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{bmatrix}^{T} = \begin{bmatrix} a_{11} & a_{21} \\ a_{12} & a_{22} \\ a_{13} & a_{23} \end{bmatrix}^{T} = \begin{bmatrix} a_{11} & a_{21} \\ a_{13} & a_{23} \end{bmatrix}^{T} = \begin{bmatrix} a_{11} & a_{21} \\ a_{21} \end{bmatrix}^{T} = \begin{bmatrix} a_{11} & a_{21} \\ a_{21} \end{bmatrix}^{T} = \begin{bmatrix} a_{11} & a_{21} \\ a_{21} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{22} & a_{23} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{22} & a_{23} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{22} & a_{23} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{22} & a_{23} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{22} & a_{23} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{22} & a_{23} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{22} & a_{23} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{22} & a_{23} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{22} & a_{23} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{23} & a_{23} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{23} & a_{23} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{23} & a_{23} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{23} & a_{23} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{23} & a_{23} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{23} & a_{23} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{23} & a_{23} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{23} & a_{23} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{23} & a_{23} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{23} & a_{23} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{23} & a_{23} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{23} & a_{23} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{23} & a_{23} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{23} & a_{23} \end{bmatrix}^{T} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{23} & a_{23} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{23} & a_{23} \end{bmatrix}^{T} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{23} & a_{23} \end{bmatrix}^{T} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{23} & a_{23} \end{bmatrix}^{T} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{23} & a_{23} \end{bmatrix}^{T} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{23} & a_{23} \end{bmatrix}^{T} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{23} & a_{23} \end{bmatrix}^{T} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{23} & a_{23} \end{bmatrix}^{T} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{23} & a_{23} \end{bmatrix}^{T} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{23} & a_{23} \end{bmatrix}^{T} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{23} & a_{23} \end{bmatrix}^{T} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{23} & a_{23} \end{bmatrix}^{T} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{23} & a_{23} \end{bmatrix}^{T} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{23} & a_{23} \end{bmatrix}^{T} \end{bmatrix}^{T} = \begin{bmatrix} a_{21} & a_{22} \\ a_{23} & a_{23} \end{bmatrix}^{T} \end{bmatrix}^{T} = \begin{bmatrix} a_{21}$$

 $-(\mathbf{A}+\mathbf{B})^{\mathrm{T}} = \mathbf{A}^{\mathrm{T}} + \mathbf{B}^{\mathrm{T}} \quad (\mathbf{A}-\mathbf{B})^{\mathrm{T}} = \mathbf{A}^{\mathrm{T}} - \mathbf{B}^{\mathrm{T}} \quad (\mathbf{A}^{\mathrm{T}})^{\mathrm{T}} = \mathbf{A} \quad (\mathbf{r}\mathbf{A})^{\mathrm{T}} = r\mathbf{A}^{\mathrm{T}}$ $-(\mathbf{A}\mathbf{B})^{\mathrm{T}} = B^{\mathrm{T}}A^{\mathrm{T}}$

6.6 Special Kinds of Matrices

A k x n matrix is a

- Square matrix if k=n that is equal number of rows and columns
 Column matrix if n=1 _____ [9]
 Row matrix if k=1 _____ [c b c f c]
- Row matrix if k=1
- Diagonal matrix if k=n and $a_{ij} = 0$ for $i \neq j$, a square matrix with nondiagonal elements are 0. If the diagonal elements are all 1 then it is called an identity matrix.
- Upper-Triangular Matrix if $a_{ij} = 0$ for i > j the entries below the dioganal is zero

e.g.
$$\begin{bmatrix} x & z \\ 0 & y \end{bmatrix}$$
, $\mathbf{L}^{\mathbf{T}} = \begin{bmatrix} 3 & 4 & 3 \\ 0 & 2 & 6 \\ 0 & 0 & 1 \end{bmatrix}$

• Lower-Triangular Matrix if $a_{ij} = 0$ for i < j the entries above the dioganal is zero

e.g.
$$\begin{bmatrix} x & 0 \\ z & y \end{bmatrix}$$
, $\mathbf{L} = \begin{bmatrix} 3 & 0 & 0 \\ 4 & 2 & 0 \\ 5 & 6 & 4 \end{bmatrix}$



• Symmetric Matrix if $A^T = A$ that is $a_{ij} = a_{jk}$

e.g.
$$\begin{bmatrix} x & z \\ z & y \end{bmatrix}$$
, $\begin{bmatrix} 5 & 6 & 4 \\ 6 & 2 & 5 \\ 4 & 5 & 1 \end{bmatrix}$, $\begin{bmatrix} 9 & 12 & 15 \\ 12 & 20 & 32 \\ 15 & 32 & 62 \end{bmatrix}$

NOTE: Any symmetric positive definite matrix ⁶ M can be written as $M=LL^{T}$ which is called Cholesky decomposition.

$$\begin{bmatrix} 9 & 12 & 15 \\ 12 & 20 & 32 \\ 15 & 32 & 62 \end{bmatrix} = \begin{bmatrix} g_{11} & 0 & 0 \\ g_{21} & g_{22} & 0 \\ g_{31} & g_{32} & g_{33} \end{bmatrix} \begin{bmatrix} g_{11} & g_{21} & g_{31} \\ 0 & g_{22} & g_{32} \\ 0 & 0 & g_{33} \end{bmatrix} = \begin{bmatrix} 3 & 0 & 0 \\ 4 & 2 & 0 \\ 5 & 6 & 1 \end{bmatrix} \begin{bmatrix} 3 & 4 & 5 \\ 0 & 2 & 6 \\ 0 & 0 & 1 \end{bmatrix}$$

Found by equating both sides $g_{11}^2 = 9 \rightarrow g_{11} = 3$ $g_{11}g_{21} = 12 \rightarrow g_{21} = 4$ $g_{11}g_{31} = 15 \rightarrow g_{31} = 5$ etc...

• Idempotent Matrix if B.B=B

e.g.
$$I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$
, $\begin{bmatrix} .5 & -5 \\ 4 & -4 \end{bmatrix} \begin{bmatrix} .5 & -5 \\ 4 & -4 \end{bmatrix} = \begin{bmatrix} 5 & -5 \\ 4 & -4 \end{bmatrix}$

• Nonsingular matrix if it is a square matrix whose rank equals the number of rows (or columns). When such a matrix is a coefficient matrix in system of linear equations, the system has one and only one solution.

2

⁶A symmetric matrix M is positive definite if the scalar z'M z is strictly positive for every nonzero column vector z

6.8 Determinant of Matrices

• Determinant is defined for square matrices. Determinant of an n x n matrix is the n-dimensional 4.1 - 3.2 = -2= 4.1 - 2.2 = 0volume scaling factor of the linear transformation produced by the matrix.

<u>__</u>_A

A.X-b

X =

• For a 2 x 2 matrix **A** its determinant is found by **Leibniz** rule:

det
$$\mathbf{A} = \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix} = a_{11}a_{22} - a_{12}a_{21}$$
 which is a scalar.

• For the determinant of 3 x 3, 4x 4 or higher size matrix, again Leibniz rule is used (multiplying the elements of a selected row or column a_{ij} with $(-1)^{i+j}$ times the *ij*th cofactor C_{ij} (determinant of the $(-1)^{\prime}(-1)^{3}=-1$ submatrix obtained by deleting row i and column j from A) and adding them up).

Note: signs come from the $(-1)^{i+j}$ where ij is the position of the first multiplicative terms (11, 12, 13, 14)for the above

$$\begin{vmatrix} a_{22} & a_{23} & a_{24} \\ a_{32} & a_{33} & a_{34} \\ a_{42} & a_{43} & a_{44} \end{vmatrix} = a_{22}(-1)^{1+1} \begin{vmatrix} a_{33} & a_{34} \\ a_{43} & a_{44} \end{vmatrix} + a_{23}(-1)^{1+2} \begin{vmatrix} a_{32} & a_{34} \\ a_{42} & a_{44} \end{vmatrix} + a_{24}(-1)^{1+3} \begin{vmatrix} a_{32} & a_{33} \\ a_{42} & a_{43} \end{vmatrix} = a_{22}a_{33}a_{44} - a_{22}a_{34}a_{43} - a_{23}a_{32}a_{44} + a_{23}a_{42}a_{34} + a_{32}a_{24}a_{43} - a_{24}a_{33}a_{42}$$

Doing for all 3 x 3 matrices and substituting the results the 4 x 4 expansion we find the determinant.

Example 44 Find the following determinant of matrix using different row or columns for the first multiplicative terms?

$$\begin{vmatrix} \mathbf{6} & -\mathbf{2} & \mathbf{2} \\ -2 & 5 & 0 \\ 2 & 0 & 7 \end{vmatrix} = 6 \begin{vmatrix} 5 & 0 \\ 0 & 7 \end{vmatrix} + 2 \begin{vmatrix} -2 & 0 \\ 2 & 7 \end{vmatrix} + 2 \begin{vmatrix} -2 & 5 \\ 2 & 7 \end{vmatrix} = 6 * 35 - 2 * 14 - 2 * 10 = 162$$

or

$$\begin{vmatrix} 6 & -2 & 2 \\ -2 & 5 & 0 \\ 2 & 0 & 7 \end{vmatrix} = 2 \begin{vmatrix} -2 & 2 \\ 5 & 0 \end{vmatrix} + 7 \begin{vmatrix} 6 & -2 \\ -2 & 5 \end{vmatrix} = -2 * 10 + 7 * 26 = 162$$
or

$$\begin{vmatrix} 6 & -2 & 2 \\ -2 & 5 & 0 \\ 2 & 0 & 7 \end{vmatrix} = 2 \begin{vmatrix} -2 & 5 \\ 2 & 0 \end{vmatrix} + 7 \begin{vmatrix} 6 & -2 \\ -2 & 5 \end{vmatrix} = -2 * 10 + 7 * 26 = 162$$

$$A^{7}_{-2} = \begin{vmatrix} 2 & 2 \\ -2 & 5 \end{vmatrix} = -2 * 10 + 7 * 26 = 162$$

$$A^{7}_{-2} = \begin{vmatrix} 2 & 2 \\ -2 & 4 \end{vmatrix} = 2 \begin{vmatrix} -2 & 2 \\ -2 & 5 \end{vmatrix} = -2 * 10 + 7 * 26 = 162$$

$$A^{7}_{-2} = \begin{vmatrix} 2 & 2 \\ -2 & 4 \end{vmatrix} = -2 * 10 + 7 * 26 = 162$$

$$A^{7}_{-2} = \begin{vmatrix} 2 & 2 \\ -2 & 4 \end{vmatrix} = -2 * 10 + 7 * 26 = 162$$

• $det(\mathbf{A}^T) = det\mathbf{A}$, $det(\mathbf{A}\mathbf{B}) = (det\mathbf{A})(det\mathbf{B})$ but $det(\mathbf{A}+\mathbf{B}) \neq (det\mathbf{A}) + (det\mathbf{B})$ in general

• A square matrix is nonsingular if and only if its determinant is nonzero.

$$A \cdot (X) = b \times - A^{-1} \cdot b$$

6.7 Inverse of Matrices

- • For a linear system of equations Ax = b we want to find the vector of unknown variables as $x = A^{-1}b$, hence we need to find the inverse of the coefficient matrix (A^{-1}) to solve the linear system easily with matrix method.
- The inverse of a matrix, A^{-1} exists only if the matrix is a square matrix. Not every square matrix has an inverse. If it has an inverse the matrix is called nonsingular, otherwise it is called singular.

tote O

- For a nonsingular matrix nxn matrix A, $AA^{-1} = I$ identity matrix
- For any nxn matrix A, let C_{ij} denote the *ij*th cofactor of A, that is, $(-1)^{i+j}$ times the determinant of the submatrix obtained by deleting row i and column j from A. The transpose of the cofactor matrix is called the **adjoint** of **A**. Then the inverse is found as:

$$A^{-1} = \frac{1}{\det A} a dj A$$

Example 45 Find the inverse of the following matrix **A**.

$$\mathbf{A} = \begin{bmatrix} 1 & -1 & 1 \\ 2 & 1 & 2 \\ 1 & 2 & 3 \end{bmatrix} \rightarrow \begin{bmatrix} -1 & 2 \\ 2 & 3 \\ 2 & 3 \end{bmatrix} = -7 \quad C_{12} = -\begin{vmatrix} 2 & 2 \\ 1 & 3 \\ 2 & 3 \end{vmatrix} = -4 \quad C_{13} = \begin{vmatrix} 2 & -1 \\ 1 & 2 \end{vmatrix} = 5$$
$$\det A = -7 * 1 - 4 * -1 + 5 * 1 = 2$$
$$C_{21} = -\begin{vmatrix} -1 & 1 \\ 2 & 3 \\ 2 & 3 \end{vmatrix} = 5 \quad C_{22} = +\begin{vmatrix} 1 & 1 \\ 1 & 3 \\ 2 & 2 \\ 2 & 2 \\ 3 & 3 \end{vmatrix} = 2 \quad C_{23} = -\begin{vmatrix} 1 & -1 \\ 1 & 2 \\ 2 & -1 \\ 1 & 2 \\ 2 & -1 \end{vmatrix} = -3$$
$$C_{31} = \begin{vmatrix} -1 & 1 \\ -1 & 2 \\ -1 & 2 \\ 2 & -1 \\ 2 & 2 \\ 2 & -1 \\ 2 & -$$

Then
$$\mathbf{C} = \begin{bmatrix} -7 & -4 & 5 \\ 5 & 2 & -3 \\ -1 & 0 & 1 \end{bmatrix} \rightarrow adjA = C^T = \begin{bmatrix} -7 & 5 & -1 \\ -4 & 2 & 0 \\ 5 & -3 & 1 \end{bmatrix} \rightarrow A^{-1} = \frac{1}{\det A}adjA = \frac{1}{2} \begin{bmatrix} -7 & 5 & -1 \\ -4 & 2 & 0 \\ 5 & -3 & 1 \end{bmatrix}$$

• 2x2 case is the most known case which is also derived from the Leibniz rule

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \rightarrow C = \begin{bmatrix} d & -c \\ -b & a \end{bmatrix} \rightarrow A^{-1} = \frac{1}{\det A} a dj A = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

$$A^{-1} = \int dt A = \int dt$$

6.8 Eigen Values and Eigen Vectors of a Matrix

An eigenvector or characteristic vector of a linear transformation is a non-zero vector that changes by only a scalar factor when that linear transformation is applied to it.Eigen values and vectors are very useful in many applications such as solving difference equations and studying stationary points higher dimensional functions.

 λ is an eigen value for an nxn matrix **A** if and only if there is a vector $v \neq 0$ and

$$\mathbf{A}v = \lambda v$$

Then the vector v is called the eigen vector of the matrix **A**.

Eigen values of a matrix is found by $\mathbf{A}v - \lambda v = (\mathbf{A} - \lambda I)v = 0$ where I is the identity matrix. So we want want for $v \neq 0$ $(\mathbf{A} - \lambda I)v = 0$ that means also $|\mathbf{A} - \lambda I| = 0$, from this determinant we find the characteristic polynomial of the matrix $P(\lambda)$ whose roots gives the eigen values of the matrix. Substituting each eigen value to $\mathbf{A}v = \lambda v$ we find the eigen vectors of the matrix for each eigen value.

Example 46 Find the eigen values of

$$A = \begin{bmatrix} 3 & 2 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 \\ 0 & 2 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -\lambda & 1 \\ 0 & 2 \end{bmatrix} = 0$$

$$\begin{bmatrix} 1 & -\lambda & 1 \\ 0 & 2 \end{bmatrix} = 0$$

$$(1 - \lambda) (2 - \lambda) = 0$$

and value $A \cdot U = \lambda \cdot U$ $(A - \lambda I) \cdot V = 0$

-1+ -1+ y=x² $A = \left| \begin{array}{c} a \\ c \\ d \end{array} \right| = ad - bc$ y' = 2x = 0 $\frac{y = 0}{\sqrt{2}} = \frac{y}{\sqrt{2}} = 0$ eters la-2 b/=0 values cd-2/=0 $\begin{vmatrix} A &= \\ | & -\lambda \\$ $(1-\lambda)(2-\lambda) - (-2.1) = 0$ $\frac{1}{4} = \frac{1}{4} = \frac{1$ $a x^2 + b x + c = 0$ rast -b = 162-hae

 $A = \begin{vmatrix} 2 & 2 \\ 5 & 5 & 6 \end{vmatrix} \qquad A^{-1} = \frac{1}{2} \begin{vmatrix} 6 & -2 \\ -5 & 2 \end{vmatrix}$ 12-10-2 det A = eigen $\begin{vmatrix} 2-\lambda & 2 \\ 5 & 6-\lambda \end{vmatrix} = 0$ x²t1 y=x² 75-87 20 $x^{2}+y^{2}=r^{2}$

7 Functions of Several Variables

7.1 Geometric representations of functions

When there are more than one variable in our functions, they can be understood by taking the resulting output or one of the variables constant and combining the graphs of them by changing constant term.

7.1.1 Graphs of Functions of two variables

• The graph of a circle can be written as $x^2 + y^2 = r^2$ where r is a constant radius. If we want to graph $f(x, y) = z = x^2 + y^2$, then we start to think from a constant radius say zero that means a point in the origin since x and y also becomes zero for f(x)=0. Increasing f(x) we get bigger radius circles and adding them up we obtain the following first graph. If we do not know the graph of the circle we may also simply think to assume y=0, then $z = x^2$ is a parabola where y=0 that means if we slice the graph at y=0 we have a parabola $z = x^2$ on xz plane, for y=-1 and y=1 we get the usual parabola pushed up one unit and for y=-2 and y=2 it is pushed up 4 units. Putting slices together we have the graph of the function. If we do the same for x we also get a parabolas also zy plane and same graph.







The graph of $z = y^2 - x^2$: Restricting y=0 we get the concave parabola $z = -x^2$, for y=1 and y=-1 we find one unit pushed version of the previous one $z = 1 - x^2$ and for y=-2 and y=2 we obtain the four unit pushed slice. Putting the slices together we graph the function.





The graph of $f(x, y) = y^2 - x^2$.



The graph of $z = y^2 - x^2$: Restricting y=0 we get the concave parabola $z = -x^2$, for y=1 and y=-1 we find one unit pushed version of the previous one $z = 1 - x^2$ and for y=-2 and y=2 we obtain the four unit pushed slice. Putting the slices together we graph the function.

7.1.2 Level Curves

Graphing the points that result the same function value is a more easier method to visualize 3D functions. Again looking on $f(x, y) = z = x^2 + y^2$, if we graph the points where the function is constant we get a circle, if we plot all the possible circles increasing the function value in a 2D space, we get the following graph. Note if we pull from the increasing parts (outside) we get the same 3d graph.



The level curves are useful to show isotherms which show the places with the same temperature and also hiking maps to see which part of the mountain is steep or flat. As seen in the following figure, at the point F the curves are close to each other such that there is close rises in altitude. So the point F is a very steep part of the mountain. On the other hand, point G seems rather flat and easy to climb.



7.1.3 Planar level sets in Economics

Economists use level sets to study the two fundemental functions of microeconomics-production function and utility function. Lets graph a simple version of Cobb-Dougless production function $(Q = kx^{\alpha}y^{\beta})$ with $k = \alpha = \beta = 1$ that is Q = xy where x is capital and y is labor. Taking production constant, we obtain the level sets which are called isoquants for the production function. For a constant Q (say 3) $y = \frac{3}{x}$, for $Q = 5 \rightarrow y = \frac{5}{x}$ and $Q = 10 \rightarrow y = \frac{10}{x}$ and $Q = 15 \rightarrow y = \frac{15}{x}$. As the isoquants increase the production increases. For different k, α and β similar level curves can be obtained. The same curves can also be used to understand the level curves of the utility function which are called indifference curves and they increase at the utility increases.



7.1.4 3D level sets

The same approach is used for 3D level sets by fixing the z value and graphing the function with fixed z using the technique we described in section 7.1.1. For example the graph of $z = x_3 + x_1^2 + x_2^2$ is the same as the graph in section 7.1.1 if z=0 and increasing z the value decreases as seen in the following graph.



Some level sets of $z = x_3 - x_1^2 - x_2^2$.

7.2 Stationary Points of Functions with Several Variables

$$f'(x) = 0$$

$$f(x) = 0$$
Stationary points are found by equating the first partial derivatives of a function to zero with respect to
each variables. Whether they are a local minimum, local maximum or a saddle point are checked similar to
checking the second derivative of a one variable function, by checking a matrix of second derivatives (called
Hessian matrix). The method can be used for functions with more variables also, but for simplicity we
only cover two and three variables cases here. For a function $f(x, y)$ or $g(x, y, z)$ with stationary point P_0
(found by $\nabla f(P_0) = 0$ or $\nabla g(P_0) = 0$) Hessian matrix is defined as

$$\begin{cases}
x = 0 \\
f(x) = 0 \\
f(x) = 0
\end{cases} = \begin{bmatrix}
f_{xx}(P_0) \\
f_{yx}(P_0) \\
f_{yx}(P_0)
\end{bmatrix} or g(x, y, z) H(P_0) = \begin{bmatrix}
g_{xx}(P_0) \\
g_{yx}(P_0) \\
g_{yx}(P_0) \\
g_{yx}(P_0)
\end{bmatrix} g_{yz}(P_0) \\
g_{yx}(P_0) \\
g_{yx}($$

In general Hessian matrices are symmetric namely $\underbrace{f_{yx}(P_0)}_{f_{xy}(P_0)} = \underbrace{f_{xy}(P_0)}_{f_{xy}(P_0)} = \underbrace{g_{yx}(P_0)}_{g_{xy}(P_0)} = \underbrace{g_{yx}(P_$

$$g(x,y) = \underbrace{e^{x}}_{(x^{2})} + \underbrace{xy - y^{2} + 3}_{2}$$

$$g_{y}(x,y) = \underbrace{x - 2y = 0 \rightarrow y = \frac{x}{2}}_{2}$$

$$g_{x}(x,y) = \underbrace{2xe^{x^{2}} + y = 0 \text{ using } y = \frac{x}{2} \rightarrow x(2e^{x^{2}} + \frac{1}{2}) = 0 \rightarrow x = y = 0 \rightarrow P_{0}(0,0)$$

$$g_{xx}(x,y) = 2e^{x^{2}} + 2x(2xe^{x^{2}}) = 2e^{x^{2}}(1 + 2x^{2}) \rightarrow g_{xx}(0,0) = 2$$

$$g_{yy}(x,y) = -2$$

$$g_{xy}(x,y) = g_{yx}(x,y) = 1$$

$$H(P_{0}) = \begin{bmatrix} 2 & 1 \\ 1 & -2 \end{bmatrix} \rightarrow \begin{vmatrix} 2 - \lambda & 1 \\ 1 & -2 - \lambda \end{vmatrix} = \lambda^{2} - 5 = 0 \rightarrow \lambda = \pm\sqrt{5} \text{ then } P_{0} \text{ is a saddle point}$$

$$f(x,y) = -2$$

$$(2x) \stackrel{!}{=} 2$$

$$(2y) \stackrel{!}{=} 2$$

$$f(x,y) = 0$$

$$f_{xy} = -2$$

$$(2x) \stackrel{!}{=} 2$$

$$(2y) \stackrel{!$$

Example 48 Study the stationary points of

$$\begin{array}{c} g(x,y,z) &= x^2 + y^4 + y^2 + z^3 - 2zz \\ g_x(x,y,z) &= 2x - 2z = 0 \to \underline{x} = \underline{z} \\ g_y(x,y,z) &= 4y^3 + 2y = 2y(2y^2 + 1) = 0 \to y = 0 \\ g_z(x,y,z) &= 4y^3 + 2y = 2y(2y^2 + 1) = 0 \to \underline{y} = 0 \\ g_z(x,y,z) &= 3z^2 - 2x \equiv z(3z - 2) = 0 \to \underline{z} \equiv 0 \text{ or } \to \underline{z} = 2/3 \text{ then two stationary points} \\ P_1 &= (0,0,0) P_2 = (2/3,0,2/3) \\ g_{xy}(x,y,z) &= 2 \\ g_{yy}(x,y,z) &= \frac{12y^2 + 2}{6z} \to g_{yy}(0,0,0) = 2 \\ g_{xy}(2,3,0,2/3) = 4 \\ g_{xy}(x,y,z) &= \frac{g_{xz}(x,y) = 0}{2} \\ g_{xy}(x,y,z) &= \frac{g_{xz}(x,y) = 0}{2} \\ g_{yz}(x,y,z) &= g_{xz}(x,y) = 0 \\ g_{yz}(x,y,z) &= g_{xz}(x,y) = 0 \\ f_1(2) = y = y \\ g_{yz}(x,y,z) &= \frac{2}{2} \\ g_{yz$$
$\begin{vmatrix} 2 & 0 & -2 \\ 0 & 2 & 0 \\ -2 & 0 & 0 \end{vmatrix} \begin{vmatrix} 2-\lambda & 0 & -2 \\ 0 & 2-\lambda & 0 \\ -2 & 0 & 0 \end{vmatrix} = \begin{vmatrix} 2-\lambda & 0 & -2 \\ 0 & 2-\lambda & 0 \\ -2 & 0 & -\lambda \end{vmatrix} = (-1)^{2+2} \cdot (2-\lambda) \begin{vmatrix} 2-\lambda & -2 \\ -2 & -\lambda \end{vmatrix}$ $-b \pm \int b^{2} hac \lambda_{1} \times z = \frac{c}{a} \qquad \lambda_{1} + \lambda_{2} = \frac{b}{a} \qquad \lambda_{i} = 2 \qquad$ $\begin{vmatrix} 2 & 0 & -2 \\ 0 & 2 & 0 \\ -2 & 0 \\ -2 & 0 \\ \end{vmatrix} = \begin{vmatrix} 2 - \lambda \\ -2 \\ -2 \\ \end{vmatrix} = \begin{pmatrix} 2 - \lambda \\ -2 \\ -2 \\ -2 \\ \end{vmatrix} = \begin{pmatrix} 2 - \lambda \\ -2 \\ -2 \\ -2 \\ \end{vmatrix} = \begin{pmatrix} 2 - \lambda \\ -2 \\ -2 \\ -2 \\ \end{vmatrix} = \begin{pmatrix} 2 - \lambda \\ -2 \\ -2 \\ -2 \\ -2 \\ \end{vmatrix}$ $(2-\lambda) \left(\frac{3^2-4\lambda}{-2\lambda} + 8 - 4 \right) - (2-\lambda) \left(\frac{\lambda^2-6\lambda}{-4\lambda} + 4 \right)$ $\lambda_1 = 2 \qquad \lambda_2 = 3 = 5 \qquad \lambda_2 = 3 = 5 \qquad \lambda_2 = 3 = 5 \qquad \lambda_3 =$ $P_2(\frac{2}{3}, 0, \frac{2}{3}) \rightarrow minimum$ all the eigen values at this point / point

axet らメ $\lambda^2 - 6\lambda + 4$ ac 36-4.1.4 6 ± 520 - 56.5 -255 6 ± 255 J5 2

Any Questions?

